

SOLUTIONS TO THE XXX TYPE BETHE ANSATZ EQUATIONS AND FLAG VARIETIES

E. MUKHIN AND A. VARCHENKO

ABSTRACT. We consider a version of the A_N Bethe equation of XXX type and introduce a reproduction procedure constructing new solutions of this equation from a given one. The set of all solutions obtained from a given one is called a population.

We show that a population is isomorphic to the sl_{N+1} flag variety and that the populations are in one-to-one correspondence with intersection points of suitable Schubert cycles in a Grassmanian variety.

We also obtain similar results for the root systems B_N and C_N . Populations of B_N and C_N type are isomorphic to the flag varieties of C_N and B_N types respectively.

1. INTRODUCTION

In this paper we consider a system of algebraic equations, see below (3.2), which we consider as an A_N version of the XXX Bethe equation. In the simplest case of $N = 1$, our system does coincide with the famous and much studied Bethe equation for the inhomogeneous XXX-model see [BIK], [Fd], [FT] and references therein.

We call solutions of our system (3.2) with additional simple conditions h -critical points and study the problem of counting the number of the h -critical points.

The definition of the critical points depends on the complex non-zero step h , complex distinct spectral parameters $z_1, \dots, z_n$, sl_N dominant integral weights $\Lambda_1, \dots, \Lambda_n$ and another sl_N weight Λ_∞ . We conjecture that for generic positions of z_i and dominant integral Λ_∞ the number of h -critical points modulo obvious symmetries equals the multiplicity of L_{Λ_∞} in $L_{\Lambda_1} \otimes \dots \otimes L_{\Lambda_n}$, where L_Λ is the irreducible sl_{N+1} representation of highest weight Λ .

In the present paper we propose a way to attack this conjecture. Given an h -critical point with a given weight at infinity Λ_∞ , we describe a procedure of constructing a set of h -critical points with weights at infinity of the form $w \cdot \Lambda_\infty$, where w is in the sl_{N+1} Weyl group and the dot denotes the shifted action, see (4.1). We call this procedure the reproduction procedure and the resulting set of h -critical points a population.

The reproduction procedure makes use of a reformulation of the algebraic system (3.2) in terms of difference equations of the second order, and furthermore in the following fertility property. Under some technical conditions, zeroes of an N -tuple of

Research of E.M. is supported in part by NSF grant DMS-0140460. Research of A.V. is supported in part by NSF grant DMS-9801582.

polynomials $(y_1, \dots, y_N)$ form an h -critical point if and only if there exist polynomials $\tilde{y}_1, \dots, \tilde{y}_N$ such that

$$W(y_i(x), \tilde{y}_i(x)) = y_{i-1}(x+h)y_{i+1}(x)T_i(x),$$

where $W(u, v) = u(x+h)v(x) - u(x)v(x+h)$ is the discrete Wronskian and polynomials T_i are given explicitly in terms of z_i and Λ_i , see (3.4). Furthermore, it turns out that in this case the zeros of the tuple $(y_1, \dots, \tilde{y}_i, \dots, y_N)$ also form an h -critical point and therefore we are able to repeat the same argument. Thus, we get a family h -critical points, each is represented by an N -tuple of polynomials.

The space V of the first coordinates of these N -tuples has dimension $N+1$ and it is called the fundamental space. The population is identified with the variety of all full flags in V . Given a flag $\{0 = F_0 \subset F_1 \subset \dots \subset F_{N+1} = V\}$, the corresponding N -tuple of polynomials is given by $y_i = W_i(F_i)/U_i$, where U_i are some explicit polynomials written in terms of T_i , see Lemma 4.9, and W_i denotes the discrete Wronskian of order i .

We consider the fundamental space of a population as an $(N+1)$ -dimensional subspace of the space $\mathbb{C}_d[x]$ of polynomials of sufficiently big degree. Therefore a fundamental space defines a point in the Grassmannian variety of $(N+1)$ -dimensional subspaces of $\mathbb{C}_d[x]$. The fact that Wronskians of all i -dimensional subspaces in V are divisible by U_i results in the conclusion that V belongs to the intersection of suitable Schubert cells in the Grassmannian. These Schubert cells are related to special flags $\mathcal{F}(z_i)$ in $\mathbb{C}_d[x]$. These flags are formed by the subspaces $F_j(z_i)$ in $\mathbb{C}_d[x]$ which consist of all polynomials divisible by $(x - z_i)(x - z_i - h) \dots (x - z_i - (d-j)h)$. In addition, the point of the Grassmannian V belongs to a specific Schubert cell related to the flag $\mathcal{F}(\infty)$. The flag $\mathcal{F}(\infty)$ is formed by the subspaces $F_j(\infty)$ in $\mathbb{C}_d[x]$ which consist of all polynomials of degree less than j .

Any population contains at most one h -critical point associated to an integral dominant weight at infinity. The fundamental spaces corresponding to different populations are different. Therefore we related the problem of counting the h -critical points associated to integral dominant weights at infinity to the problem of counting the points in intersections of Schubert cycles. It is well-known that these Schubert cycles have the algebraic index of intersection equal to the multiplicity of L_{Λ_∞} in $L_{\Lambda_1} \otimes \dots \otimes L_{\Lambda_n}$. It is an open question if for generic z_i the intersection points are all of multiplicity one.

The scheme described above repeats the method in [MV1], where a similar picture was developed for the case of Bethe equations related to the Goden model. The paper [MV1] is a smooth version of the present paper; it uses Fuchsian differential equations and the usual Wronskians.

In the smooth case, the critical points, reproduction procedure and populations are defined for any Kac-Moody algebra. It is not clear how to achieve such level of generality in the discrete situation of the present paper. We suggest the relevant definitions in the case of root systems of type B_N and C_N .

We say that zeroes of polynomials $(y_1, \dots, y_N)$ constitute a B_N (resp. C_N) h -critical point if zeroes of

$$(y_1(x), \dots, y_N(x), y_{N-1}(x+h), y_{N-2}(x+2h), \dots, y_1(x+(N-1)h))$$

(resp.

$$(y_1(x), \dots, y_N(x), y_N(x+h/2), y_{N-1}(x+3h/2), \dots, y_1(x+N-h/2)))$$

form an sl_{2N} (resp. sl_{2N+1}) h -critical point, see Section 6.1 (resp. Section 7.1) for details. Our definition is motivated by similar properties of critical points in the smooth case, see [MV1]. It turns out that h -critical points defined in such a way are exactly solutions of some systems of algebraic equations in both cases of B_N and C_N . We describe these algebraic equations.

A space of polynomials V of dimension $N+1$ is called h -selfdual if the space of discrete Wronskians $W(F)/U$, where F runs over N -dimensional subspaces of V and U is the greatest common divisor of all $W(F)$, coincides with the space of functions of the form $v(x-(N-1)h/2)$, where $v \in V$. Such a space V has a natural non-degenerate form which turns out to be symmetric if the dimension of V is odd and skew-symmetric if the dimension of V is even.

A population of B_N (resp. C_N) type is then naturally identified with the space of isotropic flags in an h -selfdual space of dimension $2N$ (resp. $2N+1$). In particular a B_N (resp. C_N) population is identified with the C_N (resp. B_N) flag variety.

The present paper deals with the h -analysis and additive shifts. Similar results can be obtained in the case of the q -analysis with multiplicative shifts related to the Bethe equations of the XXZ type.

The paper is constructed as follows. We start with the case of sl_2 in Section 2. The Sections 3 and 4 are devoted to the case of sl_{N+1} . We discuss h -selfdual spaces of polynomials in Section 5 and then deal with the cases of B_N and C_N in Sections 6 and 7. Appendix A describes in detail the simplest example of an sl_3 and “ C_1 ” populations. Appendix B collects identities involving discrete Wronskians.

We thank V. Tarasov for interesting discussions.

2. THE CASE OF sl_2

2.1. The Bethe equation. Our parameters are distinct complex numbers $z_1, \dots, z_n$, positive integers $\Lambda_1, \dots, \Lambda_n$ and a complex non-zero number h . We call z_i the *ramification points*, Λ_i the *weights* and h the *step*.

For given $l \in \mathbb{Z}_{\geq 0}$, the sl_2 *Bethe equation* is the following system of algebraic equations for the complex variables $\mathbf{t} = (t_1, \dots, t_l)$

$$\prod_{s=1}^n \frac{t_j - z_s + \Lambda_s h}{t_j - z_s} \prod_{k \neq j} \frac{t_j - t_k - h}{t_j - t_k + h} = 1, \quad (2.1)$$

where $j = 1, \dots, l$. If all Λ_i are equal to 1 then the system (2.1) is the Bethe equation for the inhomogeneous XXX model, [BIK].

Consider the product of Euler gamma functions

$$\Phi = \prod_{j=1}^l \prod_{s=1}^n \frac{\Gamma((t_j - z_s + \Lambda_s h)/p)}{\Gamma((t_j - z_s)/p)} \prod_{j,k; j < k} \frac{\Gamma((t_j - t_k - h)/p)}{\Gamma((t_j - t_k + h)/p)}.$$

The function Φ is called the master function associated to the Yangian of sl_2 and $\mathbf{z}, \mathbf{\Lambda}$. It is used in the construction of integral solutions to the rational quantum Knizhnik-Zamolodchikov equation, see [TV1], [MV2]. The function Φ is the sl_2 difference counterpart of the master function (2.1) in [MV1]. Equation (2.1) can be rewritten in the form

$$\lim_{p \rightarrow 0} \frac{\Phi(t_1, \dots, t_j + p, \dots, t_l)}{\Phi(t_1, \dots, t_j, \dots, t_l)} = 1, \quad j = 1, \dots, l.$$

If $\mathbf{t} = (t_1, \dots, t_l)$ is a solution of (2.1) then any permutation of the coordinates is also a solution. An S_l orbit of solutions of equation (2.1) such that $t_i \neq t_k$ and $t_i \neq z_s - jh$ for all $i, s, k, i \neq k$, and $j = 1, \dots, \Lambda_s$ is called an *h-critical point*.

Not all solutions of the Bethe equation are *h-critical points*. For instance, if $n = 0$ and l is even, then $t_1 = \dots = t_l = 0$ is a solution of the Bethe equation which is not an *h-critical point*.

2.2. Second order difference equations. For each *h-critical point* $\mathbf{t}$ we write the monic polynomial $u(x) = \prod_{i=1}^l (x - t_i)$ and say that a polynomial $cu(x)$ for any non-zero complex number c , represents $\mathbf{t}$. In this section we rewrite the Bethe equation in terms of a difference equation for $u(x)$.

For two functions $u(x), v(x)$, define the discrete Wronskian $W(u, v)(x)$ by the formula

$$W(u, v)(x) = u(x + h)v(x) - u(x)v(x + h).$$

Let $A(x), B(x), C(x)$ be given functions. Consider the difference equation

$$A(x) u(x + h) + B(x)u(x) + C(x)u(x - h) = 0 \tag{2.2}$$

with respect to the unknown function $u(x)$. Note that the complex vector space of polynomial solutions of equation (2.2) has dimension at most 2.

The following lemma is straightforward.

Lemma 2.1. *If $u(x)$ and $v(x)$ are two solutions of equation (2.2), then the discrete Wronskian $W(u, v)(x)$ satisfies the first order difference equation*

$$W(u, v)(x + h) = \frac{C(x + h)}{A(x + h)} W(u, v)(x). \tag{2.3}$$

□

Given ramification points z_i and weights Λ_i we now fix the choice of the functions $A(x)$ and $C(x)$ as follows

$$A(x) = \prod_{s=1}^n (x - z_s), \quad C(x) = \prod_{s=1}^n (x - z_s + \Lambda_s h). \quad (2.4)$$

Then the polynomial solutions of equation (2.3) are constant multiples of the function

$$T(x) = \prod_{s=1}^n \prod_{i=1}^{\Lambda_s} (x - z_s + ih). \quad (2.5)$$

We say that a polynomial $u(x)$ is *generic* with respect to $\mathbf{z} = (z_1, \dots, z_n)$, $\mathbf{\Lambda} = (\Lambda_1, \dots, \Lambda_n)$ if the polynomial $u(x)$ has no multiple roots and no common roots with polynomials $u(x+h)$, $T(x)$.

Lemma 2.2. *Assume that $B(x)$ is a polynomial and that a solution $u(x)$ of (2.2) is a polynomial generic with respect to $\mathbf{z}, \mathbf{\Lambda}$. Then $u(x)$ represents an h -critical point.*

Conversely, if $u(x)$ represents an h -critical point then the polynomial $u(x)$ satisfies equation (2.2), where $B(x)$ is a polynomial given by

$$B(x) = - \left(\prod_{s=1}^n (x - z_s) u(x+h) + \prod_{s=1}^n (x - z_s + \Lambda_s h) u(x-h) \right) / u(x). \quad (2.6)$$

Proof. The lemma follows from the fact that equation (2.1) with respect to t_j is obtained by the substituting t_j in equation (2.2). $\square$

2.3. The sl_2 reproduction. By Lemma 2.2, each h -critical point defines a difference equation together with a polynomial solution. This difference equation is a linear difference equation of the second order with polynomial coefficients. In this section we show that the space of solutions of that difference equation contains a complex two-dimensional space of polynomials all of which (except for scalar multiples of finitely many) represent h -critical points.

Lemma 2.3. *Let the roots of the polynomial $u(x)$ be distinct and satisfy the Bethe equation (2.1). Then the corresponding difference equation (2.2) has two linearly independent polynomial solutions.*

Proof. Let $u(x) = \prod_{i=1}^n (x - t_i)$. We look for the second polynomial solution $v(x)$ in the form $v(x) = c(x)u(x)$. After multiplication by a non-zero number, we have the equation

$$v(x+h)u(x) - v(x)u(x+h) = \prod_{s=1}^n \prod_{i=1}^{\Lambda_s} (x - z_s + ih). \quad (2.7)$$

Therefore $c(x)$ is a solution of the equation

$$c(x+h) - c(x) = \frac{\prod_{s=1}^n \prod_{i=1}^{\Lambda_s} (x - z_s + ih)}{u(x)u(x+h)}. \quad (2.8)$$

The right hand side of this equation is of the form

$$f(x) + \sum_{j=1}^l \left(\frac{a_j}{x - t_j + h} - \frac{b_j}{x - t_j} \right),$$

where $f(x)$ is a polynomial and a_j, b_j are some numbers. Moreover, equation (2.1) imply that $a_j = b_j$ for all j .

The equation $c(x+h) - c(x) = f(x)$ has a polynomial solution. Denote it $\tilde{c}(x)$. Then $c(x) = \tilde{c}(x) + \sum_{j=1}^l a_j/(x - t_j)$ is a solution to (2.8).

Then the function $v(x) = c(x)u(x)$ is a polynomial solution of equation (2.2). $\square$

Thus, starting from a solution $\mathbf{t}$ of the Bethe equation (2.1), we obtain a difference equation of order two which has a two-dimensional space of polynomial solutions. We call the projectivisation of this space the *population of h -critical points related to $\mathbf{t}$* and denote $P(\mathbf{t})$.

Generic polynomials with respect to $\mathbf{z}, \mathbf{\Lambda}$ form a Zariski open subset of the population. Roots of a generic polynomial form an h -critical point by Lemma 2.2.

Let $\partial : f(x) \rightarrow f(x+h)$ be the shift operator acting on functions of one variable. The difference operator $\partial^2 + B(x)/A(x)\partial + C(x)/A(x)$ obtained from an h -critical point $\bar{\mathbf{t}} \in P(\mathbf{t})$ does not depend on the choice of $\bar{\mathbf{t}}$. We call this operator the *fundamental operator associated to the population $P(\mathbf{t})$* and denote it by $D_{P(\mathbf{t})}$.

2.4. Fertile polynomials. We give another equivalent condition for a polynomial to define an h -critical point.

Lemma 2.4. *Let a polynomial $u(x) = \prod_{i=1}^l (x - t_i)$ be generic with respect to $\mathbf{z}, \mathbf{\Lambda}$. Then the roots of $u(x)$ form a solution of equation (2.1) if and only if there exists a polynomial solution $v(x)$ to equation (2.7).*

Proof. If the roots of $u(x)$ form a solution of equation (2.5), then equation (2.2) with the polynomial $B(x)$ given by (2.6) has two linearly independent polynomial solutions by Lemma 2.3, and hence equation (2.7) has a polynomial solution.

Now assume that $v(x)$ is a polynomial solution to (2.7). Then the rational function $c(x) = v(x)/u(x)$ is a solution of equation (2.8). Then

$$\operatorname{Res}_{x=t_j} \frac{\prod_{s=1}^n \prod_{i=1}^{\Lambda_s} (x - z_s + ih)}{u(x)u(x+h)} + \operatorname{Res}_{x=t_j-h} \frac{\prod_{s=1}^n \prod_{i=1}^{\Lambda_s} (x - z_s + ih)}{u(x)u(x+h)} = 0$$

for $j = 1, \dots, l$. This system of equations is equivalent to equation (2.1). $\square$

A polynomial $u(x)$ is called *fertile* if there exist a polynomial $v(x)$ such that the discrete Wronskian of u and v is $T(x)$, $W(u, v)(x) = T(x)$. All polynomials in a population of h -critical points are fertile. Moreover, the h -critical points correspond to fertile generic polynomials.

We get the following immediate corollary.

Lemma 2.5. *If equation (2.2) has two polynomial solutions of degree l and l' , $l \neq l'$, then $l + l' - 1 = \Lambda_1 + \dots + \Lambda_n$. $\square$*

3. THE CASE OF sl_{N+1}

Let $\alpha_1, \dots, \alpha_N$ be the simple roots of sl_{N+1} . We have $(\alpha_i, \alpha_i) = 2$, $(\alpha_i, \alpha_{i\pm 1}) = -1$ with all other scalar products equal to zero.

3.1. Definition of h -critical points. In the case of sl_{N+1} we fix the following parameters: *ramification points* $\mathbf{z} = (z_1, \dots, z_n) \in \mathbb{C}^n$; non-zero dominant integral sl_{N+1} *weights* $\mathbf{\Lambda} = (\Lambda_1, \dots, \Lambda_n)$, *relative shifts* $\mathbf{b}^{(i)} = (b_1^{(i)}, \dots, b_n^{(i)}) \in \mathbb{C}^n$, $i = 1, \dots, N$, and the step $h \in \mathbb{C}$, $h \neq 0$.

We call the set of parameters $\mathbf{z}, \mathbf{\Lambda}, \mathbf{b}^{(i)}$, $i = 1, \dots, N$, the *initial data*. We denote $\Lambda_s^{(i)} = (\Lambda_s, \alpha_i) \in \mathbb{Z}_{\geq 0}$.

Let $\mathbf{l} = (l_1, \dots, l_N) \in \mathbb{Z}_{\geq 0}^N$. The sl_{N+1} -weight

$$\Lambda_\infty = \sum_{s=1}^n \Lambda_s - \sum_{i=1}^N l_i \alpha_i, \quad (3.1)$$

is called the *weight at infinity*.

The following system of algebraic equations for variables $\mathbf{t} = (t_j^{(i)})$, $i = 1, \dots, N$, $j = 1, \dots, l_i$,

$$\prod_{s=1}^n \frac{t_j^{(i)} - z_s + b_s^{(i)} h + \Lambda_s^{(i)} h}{t_j^{(i)} - z_s + b_s^{(i)} h} \prod_{k=1}^{l_{i-1}} \frac{t_j^{(i)} - t_k^{(i-1)} + h}{t_j^{(i)} - t_k^{(i-1)}} \prod_{k \neq j} \frac{t_j^{(i)} - t_k^{(i)} - h}{t_j^{(i)} - t_k^{(i)} + h} \times \prod_{k=1}^{l_{i+1}} \frac{t_j^{(i)} - t_k^{(i+1)}}{t_j^{(i)} - t_k^{(i+1)} - h} = 1, \quad (3.2)$$

is called the sl_{N+1} *Bethe equation* associated with the initial data and the weight at infinity.

Note that by a shift of variables $\tilde{t}_i^{(j)} = t_i^{(j)} - jh/2$ the system can be written in a slightly more symmetrical way.

Note that in the quasiclassical limit $h \rightarrow 0$, system (3.2) becomes system (2.2) of [MV1] specialized to the case of sl_{N+1} .

The product of symmetric groups $S_{\mathbf{l}} = S_{l_1} \times \dots \times S_{l_N}$ acts on the set of solutions of (3.2) permuting the coordinates with the same upper index. An $S_{\mathbf{l}}$ orbit of solutions of the Bethe equation such that $t_j^{(i)} \neq t_k^{(i)}$, $t_j^{(i)} \neq z_s - b_s^{(i)} - rh$ for all j, i, s, k , $k \neq j$

and $r = 1, \dots, \Lambda_s^{(i)}$ is called an *h-critical point associated with the initial data and the weight at infinity*.

Let L_Λ be the irreducible sl_{N+1} module with highest weight Λ . Let the initial data satisfies $b_s^{(j)} = -\sum_{i=1}^j \Lambda_s^{(i)}$. We have the following conjecture.

Conjecture 3.1. *If Λ_∞ is integral dominant then for generic $\mathbf{z}$ the number of h-critical points associated with the initial data and the weight at infinity equals to the multiplicity of L_{Λ_∞} in $L_{\Lambda_1} \otimes \dots \otimes L_{\Lambda_n}$.*

3.2. Difference equations of the second order. In this section, given an *h-critical point* we obtain N difference operators of order two. The i th operator is the fundamental operator with respect to the $sl_2 \in sl_{N+1}$ in the direction α_i .

Let $\mathbf{t}$ be an *h-critical point*. For $i = 1, \dots, N$, introduce polynomials

$$y_i(x) = \prod_{j=1}^{l_i} (x - t_j^{(i)}). \quad (3.3)$$

We also set $y_0(x) = 1$, $y_{N+1}(x) = 1$.

The N -tuple $\mathbf{y}$ uniquely determines the *h-critical point* $\mathbf{t}$ and we say that $\mathbf{y}$ *represents* $\mathbf{t}$. We consider the tuple $\mathbf{y}$ up to multiplication of each coordinate by a non-zero number, since we are interested only in the roots of polynomials $y_1, \dots, y_N$. Thus the tuple defines a point in the direct product $\mathbf{P}(\mathbb{C}[x])^N$ of N copies of the projective space associated with the vector space of polynomials of x . In this paper we view all N -tuples of polynomials as elements of $\mathbf{P}(\mathbb{C}[x])^N$.

Introduce the polynomials

$$\begin{aligned} T_i(x) &= \prod_{s=1}^n \prod_{j=1}^{\Lambda_s^{(i)}} (x - z_s + b_s^{(i)}h + jh), \\ A_i(x) &= \prod_{s=1}^n (x - z_s + b_s^{(i)}h) \prod_{k=1}^{l_{i-1}} (x - t_k^{(i-1)}) \prod_{k=1}^{l_{i+1}} (x - t_k^{(i+1)} - h), \\ C_i(x) &= \prod_{s=1}^n (x - z_s + b_s^{(i)}h + \Lambda_s^{(i)}h) \prod_{k=1}^{l_{i-1}} (x - t_k^{(i-1)} + h) \prod_{k=1}^{l_{i+1}} (x - t_k^{(i+1)}). \end{aligned} \quad (3.4)$$

We say that a tuple of polynomials $\mathbf{y} \in \mathbf{P}(\mathbb{C}[x])^N$ is *generic with respect to the initial data* if for all i the polynomial $y_i(x)$ has no multiple roots and no common roots with polynomials $y_i(x+h)$, $y_{i-1}(x+h)$, $y_{i+1}(x)$, $T_i(x)$.

For $i = 1, \dots, N$, consider difference equations of the second order of the form

$$A_i(x)u(x+h) + B_i(x)u(x) + C_i(x)u(x-h) = 0, \quad (3.5)$$

where $B_i(x)$ are any functions.

Lemma 3.2. *Assume that for all i , y_i is a polynomial solution of i -th equation in (3.5) for some polynomial $B_i(x)$. If the tuple $\mathbf{y} = (y_1, \dots, y_N)$ is generic with respect to the initial data then $\mathbf{y}$ represents an h -critical point.*

Conversely, if a tuple of polynomials $\mathbf{y} = (y_1, \dots, y_N)$ represents an h -critical point, then for all i the polynomial y_i satisfies the i -th equation in (3.5), where $B_i(x)$ is a polynomial given by $B_i(x) = -(A_i(x)y_i(x+h) + C_i(x)y_i(x-h))/y_i(x)$.

Proof. The lemma follows directly from the sl_2 counterpart, Lemma 2.2. $\square$

3.3. Fertile tuples. In this section we discuss yet another criteria for a tuple of polynomials to represent an h -critical point.

We say that a tuple of polynomials $\mathbf{y} \in \mathbf{P}(\mathbb{C}[x])^N$ is *fertile with respect to the initial data* if for every i there exists a polynomial $\tilde{y}_i$ such that the discrete Wronskian is

$$W(y_i, \tilde{y}_i)(x) = T_i(x)y_{i-1}(x+h)y_{i+1}(x). \quad (3.6)$$

Lemma 3.3. *A generic tuple $\mathbf{y}$ is fertile if and only if it represents an h -critical point.*

Proof. The lemma follows from the sl_2 considerations, see Lemma 2.4. $\square$

Let $\mathbf{y}$ be fertile and let $y_i, \tilde{y}_i$ satisfy (3.6). Then polynomials of the form $c_1 y_i + c_2 \tilde{y}_i$ with $c_1, c_2 \in \mathbb{C}$ span the two dimensional space of polynomial solutions of equation (3.5). The tuples $(y_1, \dots, c_1 y_i + c_2 \tilde{y}_i, \dots, y_N) \in \mathbf{P}(\mathbb{C}[x])^N$ are called *immediate descendants* of $\mathbf{y}$ in the i -th direction. Note that if $\mathbf{y}$ is generic, then all but finitely many immediate descendants are generic.

Let $\mathbb{C}_d[x]$ be the space of polynomials of degree not greater than d . The set of fertile tuples is closed in $\mathbf{P}(\mathbb{C}_d[x])^N$:

Lemma 3.4. *Assume that a sequence of fertile tuples of polynomials $\mathbf{y}_k$, $k = 1, 2, \dots$, has a limit $\mathbf{y}_\infty$ in $\mathbf{P}(\mathbb{C}_d[x])^N$ as k tends to infinity. Then the limiting tuple $\mathbf{y}_\infty$ is fertile. Moreover, if $\mathbf{y}_\infty^{(i)}$ is an immediate descendant of $\mathbf{y}_\infty$ in the i -th direction, then there exist immediate descendants $\mathbf{y}_k^{(i)}$ of $\mathbf{y}_k$ in the i -th direction such that $\mathbf{y}_\infty^{(i)}$ is the limit of $\mathbf{y}_k^{(i)}$.*

Proof. Let us prove $\mathbf{y}_\infty$ is fertile in direction i . For each k we have a difference equation (3.5) which we denote E_k and a plane of polynomial solutions P_k . By the assumptions of the lemma there is a limiting equation E_∞ as $k \rightarrow \infty$.

The space P_k defines a point in the projective Grassmanian variety of planes in $\mathbb{C}_d[x]^N$. Let P_∞ be a limiting plane, then all polynomials in P_∞ are solutions of equation E_∞ and the lemma is proved. $\square$

3.4. Reproduction procedure. Given an h -critical point, we describe a procedure of obtaining a family of h -critical points.

Theorem 3.5. *Let $\mathbf{y}$ represent an h -critical point and let $y_i, \tilde{y}_i$ satisfy (3.6). Assume that $\mathbf{y}^{(i)} = (y_1, \dots, \tilde{y}_i, \dots, y_N)$ is generic. Then $\mathbf{y}^{(i)}$ represents an h -critical point.*

Proof. Denote $\bar{t}_j^{(i)}$ the roots of $\bar{y}_i = c_1 y_i + c_2 \tilde{y}_i$. Let z be a root of $T_i(x) y_{i-1}(x+h) y_{i+1}(x)$. Then we have

$$\frac{y_i(z+h)}{y_i(z)} = \frac{\bar{y}_i(z+h)}{\bar{y}_i(z)}.$$

In particular we choose $z = t_k^{(i-1)} - h$ and $z = t_k^{(i+1)}$ and obtain

$$\begin{aligned} \prod_j \frac{t_k^{(i-1)} - t_j^{(i)}}{t_k^{(i-1)} - t_j^{(i)} - h} &= \prod_j \frac{t_k^{(i-1)} - \bar{t}_j^{(i)}}{t_k^{(i-1)} - \bar{t}_j^{(i)} - h}, \\ \prod_j \frac{t_k^{(i+1)} - t_j^{(i)} + h}{t_k^{(i+1)} - t_j^{(i)}} &= \prod_j \frac{t_k^{(i+1)} - \bar{t}_j^{(i)} + h}{t_k^{(i+1)} - \bar{t}_j^{(i)}} \end{aligned}$$

for all k and i . In addition Lemma 2.2 implies that for all k ,

$$\prod_{j; j \neq k} \frac{t_k^{(i)} - t_j^{(i)} - h}{t_k^{(i)} - t_j^{(i)} + h} = \prod_{j; j \neq k} \frac{\bar{t}_k^{(i)} - \bar{t}_j^{(i)} - h}{\bar{t}_k^{(i)} - \bar{t}_j^{(i)} + h}.$$

That proves the theorem. $\square$

Thus, starting with a tuple $\mathbf{y}$, representing an h -critical point and an index $i \in \{1, \dots, N\}$, we construct a closed subvariety of immediate descendants $\mathbf{y}^{(i)}$ in $\mathbf{P}(\mathbb{C}[x])^N$ which is isomorphic to $\mathbf{P}^1$. All but finitely many points in this subvariety correspond to N -tuples of polynomials which represent h -critical points. All constructed N -tuples of polynomials are fertile. We call this construction the *simple reproduction procedure in the i -th direction*.

Now we can repeat the simple reproduction procedure in some other j -th direction applied to all points obtained in the previous step and obtain more critical points.

We continue the process until no simple reproduction in any direction applied to any polynomial obtained in the previous steps produces a new polynomial. The result is called the *population originated at the h -critical point $\mathbf{y}$* and is denoted $P(\mathbf{y})$.

More formally, the population of $\mathbf{y}$ is the set of all N -tuples of polynomials $\bar{\mathbf{y}} \in \mathbf{P}(\mathbb{C}[x])^N$ such that there exist N -tuples $\mathbf{y}_1 = \mathbf{y}$, $\mathbf{y}_2, \dots, \mathbf{y}_k = \bar{\mathbf{y}}$, such that $\mathbf{y}_i$ is an immediate descendant of $\mathbf{y}_{i-1}$ for all $i = 2, \dots, k$.

Obviously, if two populations intersect then they coincide.

4. FUNDAMENTAL SPACE OF AN sl_{N+1} POPULATION

Given an sl_{N+1} population of critical points we construct a space of polynomials called the fundamental space. Elements of the population are in natural correspondence with full flags in the fundamental space. Then we show that the problem of counting h -critical points is equivalent to a problem of Schubert calculus.

4.1. Degrees of polynomials in a population and the sl_{N+1} Weyl group. Let $\mathfrak{h}$ be the Cartan subalgebra of sl_{N+1} and let $(\cdot, \cdot)$ be the standard scalar product on $\mathfrak{h}^*$. Denote $\rho \in \mathfrak{h}^*$ the half sum of positive roots.

The Weyl group $\mathcal{W} \in \text{End}(\mathfrak{h}^*)$ is generated by reflections $s_i, i = 1, \dots, N$,

$$s_i(\lambda) = \lambda - (\lambda, \alpha_i)\alpha_i, \quad \lambda \in \mathfrak{h}^*.$$

We use the notation

$$w \cdot \lambda = w(\lambda + \rho) - \rho, \quad w \in \mathcal{W}, \lambda \in \mathfrak{h}^*, \quad (4.1)$$

for the shifted action of the Weyl group.

Let an initial data be given. Let $\mathbf{y} = (y_1, \dots, y_N) \in \mathbf{P}(\mathbb{C}[x])^N$, and let l_i be the degree of the polynomial y_i for $i = 1, \dots, N$. Let Λ_∞ be the weight at infinity defined by (3.1).

Assume the tuple $\mathbf{y}$ is fertile. Let the tuple $\mathbf{y}^{(i)} = (y_1, \dots, \tilde{y}_i, \dots, y_N)$ be an immediate descendant of $\mathbf{y}$ in the direction i and let $\Lambda_\infty^{(i)}$ be the sl_{N+1} weight at infinity of $\mathbf{y}^{(i)}$.

The following lemma follows from Lemma 2.5.

Lemma 4.1. *If the degree of $\tilde{y}_i$ is not equal to the degree of y_i , then*

$$\Lambda_\infty^{(i)} = s_i \cdot \Lambda_\infty,$$

where $s_i \cdot$ is the shifted action of the i -th generating reflection of the Weyl group.

Therefore we obtain the following

Proposition 4.2. *Let a tuple $\mathbf{y}^0$ represent an h -critical point associated with a given initial data and weight at infinity Λ_∞ . Let P be the population of h -critical points originated at $\mathbf{y}^0$. Then*

- *For any tuple $\mathbf{y} \in P$, there is an element w of the sl_{N+1} Weyl group $\mathcal{W}$, such that the weight at infinity of $\mathbf{y}$ is $w \cdot \Lambda_\infty$.*
- *For any element $w \in \mathcal{W}$, there is a tuple $\mathbf{y} \in P$ whose weight at infinity is $w \cdot \Lambda_\infty$.*

□

Proposition 4.2 gives sufficient conditions for absence of h -critical points.

Corollary 4.3. *There is no h -critical point in either of the two cases*

- (1) *if there is an element w of the Weyl group such that $\sum_{s=1}^n \Lambda_s - w \cdot \Lambda_\infty$ does not belong to the cone $\mathbb{Z}_{\geq 0}\alpha_1 \oplus \dots \oplus \mathbb{Z}_{\geq 0}\alpha_N$;*
- (2) *if Λ_∞ belongs to one of the reflection hyperplanes of the shifted action of the Weyl group.*

The next corollary says that under certain conditions on weights there is exactly one population of h -critical points.

The tuple $(1, \dots, 1) \in \mathbf{P}(\mathbb{C}[x])^N$ is the unique N -tuple of non-zero polynomials of degree 0. The weight at infinity of $(1, \dots, 1)$ is $\Lambda_{\infty, (1, \dots, 1)} = \sum_{s=1}^n \Lambda_s$. Let $P_{(1, \dots, 1)}$ be the population associated to the initial data and originated at $(1, \dots, 1)$.

Corollary 4.4. *Let $\mathbf{y}$ be an h -critical point such that Λ_∞ has the form $w \cdot \Lambda_{\infty,(1,\dots,1)}$ for some $w \in \mathcal{W}$. Then $\mathbf{y}$ belongs to the population $P_{(1,\dots,1)}$.*

4.2. The difference operator of a population. In this section we describe a linear difference operator of order $N + 1$ related to a population of h -critical points.

Let an initial data $\mathbf{z}, \mathbf{\Lambda}, \mathbf{b}^{(i)}$ be given. Let the polynomials T_i , $i = 1, \dots, N$, be defined by formula (3.4).

Let $\mathbf{y} = (y_1, \dots, y_N)$ be an N -tuple of non-zero polynomials. Set $y_0 = y_{N+1} = 1$. Define a linear difference operator of order $N + 1$ with meromorphic coefficients

$$\begin{aligned} D(\mathbf{y}) &= \left(\partial - \frac{y_N(x)}{y_N(x+h)} \prod_{s=1}^N \frac{T_s(x + (N-s+1)h)}{T_s(x + (N-s)h)} \right) \times \\ &\quad \left(\partial - \frac{y_N(x+h)}{y_N(x)} \frac{y_{N-1}(x)}{y_{N-1}(x+h)} \prod_{s=1}^{N-1} \frac{T_s(x + (N-s)h)}{T_s(x + (N-1-s)h)} \right) \times \dots \times \\ &\quad \left(\partial - \frac{y_2(x+h)}{y_2(x)} \frac{y_1(x)}{y_1(x+h)} \frac{T_1(x+h)}{T_1(x)} \right) \left(\partial - \frac{y_1(x+h)}{y_1(x)} \right) = \\ &= \prod_i^{0 \rightarrow N} \left(\partial - \frac{y_{N+1-i}(x+h)}{y_{N+1-i}(x)} \frac{y_{N-i}(x)}{y_{N-i}(x+h)} \prod_{s=1}^{N-i} \frac{T_s(x + (N-i-s+1)h)}{T_s(x + (N-i-s)h)} \right). \end{aligned} \quad (4.2)$$

Notice that the first coordinate y_1 of the N -tuple belongs to the kernel of the operator $D(\mathbf{y})$.

Theorem 4.5. *Let P be a population of h -critical points originated at some tuple $\mathbf{y}^0$. Then the operator $D(\mathbf{y})$ does not depend on the choice of $\mathbf{y}$ in P .*

Proof. We have to prove that if $\mathbf{y}, \tilde{\mathbf{y}} \in P$ then $D(\mathbf{y}) = D(\tilde{\mathbf{y}})$. It is enough to show that is the case when $\tilde{\mathbf{y}}$ is an immediate descendant of $\mathbf{y}$ in some direction i and both $\mathbf{y}, \tilde{\mathbf{y}}$ represent critical points. Therefore, we assume that $y_j = \tilde{y}_j$, for all $j, j \neq i$, and $W(y_i, \tilde{y}_i)(x) = T_i(x) y_{i-1}(x+h) y_{i+1}(x)$. In this case, all factors of $D(\mathbf{y})$ and $D(\tilde{\mathbf{y}})$ are the same except the two factors which involve y_i or $\tilde{y}_i$. So it is left to show that for

any function $u(x)$ we have

$$\begin{aligned}
& \left(\partial - \frac{y_{i+1}(x+h)}{y_{i+1}(x)} \frac{y_i(x)}{y_i(x+h)} \prod_{s=1}^i \frac{T_s(x+(i-s+1)h)}{T_s(x+(i-s)h)} \right) \times \\
& \left(\partial - \frac{y_i(x+h)}{y_i(x)} \frac{y_{i-1}(x)}{y_{i-1}(x+h)} \prod_{s=1}^{i-1} \frac{T_s(x+(i-s)h)}{T_s(x+(i-s-1)h)} \right) u(x) = \\
& \left(\partial - \frac{y_{i+1}(x+h)}{y_{i+1}(x)} \frac{\tilde{y}_i(x)}{\tilde{y}_i(x+h)} \prod_{s=1}^i \frac{T_s(x+(i-s+1)h)}{T_s(x+(i-s)h)} \right) \times \\
& \left(\partial - \frac{\tilde{y}_i(x+h)}{\tilde{y}_i(x)} \frac{y_{i-1}(x)}{y_{i-1}(x+h)} \prod_{s=1}^{i-1} \frac{T_s(x+(i-s)h)}{T_s(x+(i-s-1)h)} \right) u(x) .
\end{aligned}$$

After the change of variables

$$v(x) = u(x) \frac{\prod_{s=1}^{i-1} T_s(x+(i-s-1)h)}{y_{i-1}(x)},$$

we have to prove that

$$\begin{aligned}
& \left(\partial - \frac{y_{i+1}(x+h)}{y_{i+1}(x)} \frac{y_i(x)}{y_i(x+h)} \frac{y_{i-1}(x+2h)}{y_{i-1}(x+h)} \frac{T_i(x+h)}{T_i(x)} \right) \left(\partial - \frac{y_i(x+h)}{y_i(x)} \right) v(x) = \\
& \left(\partial - \frac{y_{i+1}(x+h)}{y_{i+1}(x)} \frac{\tilde{y}_i(x)}{\tilde{y}_i(x+h)} \frac{y_{i-1}(x+2h)}{y_{i-1}(x+h)} \frac{T_i(x+h)}{T_i(x)} \right) \left(\partial - \frac{\tilde{y}_i(x+h)}{\tilde{y}_i(x)} \right) v(x) .
\end{aligned}$$

This identity is easily checked directly using the equation connecting y_i and $\tilde{y}_i$. $\square$

The difference operator (4.2) is called the *fundamental operator associated to the population P* and is denoted D_P .

Corollary 4.6. *Let $\mathbf{y}$ be a member of a population P . Then the first coordinate y_1 of $\mathbf{y}$ is in the kernel of the operator D_P .*

Proposition 4.7. *The space of polynomial solutions to the difference equation*

$$D_P u = 0 \tag{4.3}$$

has dimension $N+1$.

Proof. Let $\mathbf{y}$ be a member of the population P . Assume that $\mathbf{y}$ represents an h -critical point. We construct polynomials $u_1, \dots, u_{N+1}$, satisfying equation (4.3).

Set $u_1 = y_1$. The polynomial u_1 is a solution of (4.3).

Let u_2 be a polynomial such that $W(u_1, u_2)(x) = T_1(x)y_2(x)$. The polynomial u_2 is a solution of (4.3).

Let $\tilde{y}_2$ be a polynomial such that $W(y_2, \tilde{y}_2)(x) = T_2(x)y_1(x+h)y_3(x)$. Such a polynomial can be chosen so that $(y_1, \tilde{y}_2, \dots, y_N)$ is generic and therefore represents an h -critical point. Choose a polynomial u_3 to satisfy equation $W(u_3, y_1)(x) = T_1(x)\tilde{y}_2(x)$. The polynomial u_3 is a solution of (4.3).

In general, to construct a polynomial u_{i+1} we find $\tilde{\mathbf{y}}_i = (y_1, \dots, \tilde{y}_i, \dots, y_N)$, such that $\tilde{\mathbf{y}}_i$ is generic and $W(y_i, \tilde{y}_i)(x) = T_i(x)y_{i-1}(x+h)y_{i+1}(x)$ and then repeat the construction for u_i using $\tilde{\mathbf{y}}_i$ instead of $\mathbf{y}$.

Let V be the complex vector space spanned by polynomials $u_1, \dots, u_{N+1}$. We show that the space V has dimension $N+1$.

Let $W(g_1, \dots, g_s)$ be the discrete Wronskian of functions $g_1, \dots, g_s$ defined by

$$W(g_1, \dots, g_s)(x) = \det(g_i(x + (j-1)h))_{i,j=1}^s$$

For $s = 0$, we define the corresponding discrete Wronskian to be 1.

Lemma 4.8. *For $i = 1, \dots, N+1$, we have*

$$W(u_1, \dots, u_i)(x) = y_i(x) \prod_{j=1}^{i-1} T_1(x + (j-1)h) \prod_{j=1}^{i-2} T_2(x + (j-1)h) \dots T_{i-1}(x).$$

Proof. The lemma is proved in the same way as Lemma 5.5 in [MV1] making use of Wronskian identities in Lemmas 9.3 and 9.5. $\square$

The proposition is proved. $\square$

A linear difference equation of order $N+1$ cannot have more than $N+1$ polynomial solutions linearly independent over $\mathbb{C}$. The complex $(N+1)$ -dimensional vector space of polynomial solutions of equation (4.3) is called *the fundamental space of the population* P and is denoted V_P .

4.3. Frames. We describe an h -analogue of ramification properties of a space of polynomials.

A space of polynomials V is called a *space without base points* if for any $z \in \mathbb{C}$ there exists $v \in V$ such that $v(z) \neq 0$.

Let V be an $(N+1)$ -dimensional vector space of polynomials without base points. Let $U_i(x)$ be the monic polynomial which is the greatest common divisor of the family of polynomials $\{W(u_1, \dots, u_i) \mid u_1, \dots, u_i \in V\}$.

Lemma 4.9. *There exist a unique sequence of monic polynomials $T_1(x), \dots, T_N(x)$ such that*

$$U_i(x) = \prod_{j=1}^{i-1} T_1(x + (j-1)h) \prod_{j=1}^{i-2} T_2(x + (j-1)h) \dots T_{i-1}(x);$$

for $i = 1, \dots, N+1$.

We call the sequence of monic polynomials $T_1(x), \dots, T_N(x)$ the *frame* of V .

Proof. We construct the polynomials T_i by induction on i . For $i = 0$, we have $U_1 = 1$. For $i = 1$ we just set $T_1 = U_2$. Suppose the lemma is proved for all $i = 1, \dots, i_0 - 1$.

Then we set

$$S(x) = \prod_{i=1}^{i_0-2} \prod_{j=1}^{i_0-i} T_i(x + (j-1)h), \quad T_{i_0-1}(x) = U_{i_0}(x)/S(x).$$

We only have to show that T_{i_0-1} is a polynomial. In other words, we have to show that a Wronskian of any i_0 dimensional subspace in V is divisible by $S(x)$.

Consider the Grassmanian $Gr(i_0 - 2, V)$ of $(i_0 - 2)$ -dimensional spaces in V . For any $z \in \mathbb{C}$ the set of points in $Gr(i_0 - 2, V)$ such that the corresponding Wronskian divided by U_{i_0-2} does not vanish at z , is a Zariski open algebraic set. Therefore we have a Zariski open set of points in $Gr(i_0 - 2, V)$ such that the corresponding Wronskian divided by U_{i_0-2} does not vanish at roots of $S(x-h)$. We call such subspaces acceptable.

Therefore we have a Zariski open set of points in $Gr(i_0, V)$ such that the corresponding i_0 dimensional space contains an acceptable $i_0 - 2$ dimensional subspace. Let $u_1, \dots, u_{i_0} \in V$ are such that $u_1, \dots, u_{i_0-2}$ span an acceptable space. It is enough to show that $W(u_1, \dots, u_{i_0})$ is divisible by $S(x)$.

Then using Wronskian identities in Lemmas 9.3 and 9.5 we have for suitable polynomials f_1, f_2, g :

$$\begin{aligned} W(u_1, \dots, u_{i_0}) &= \frac{W(W(u_1, \dots, u_{i_0-1}), W(u_1, \dots, u_{i_0-2}, u_{i_0}))}{W(u_1, \dots, u_{i_0-2})(x+h)} = \\ &= \frac{W(U_{i_0-1}f_1, U_{i_0-1}f_2)}{U_{i_0-2}(x+h)g(x+h)} = \frac{U_{i_0-1}(x)U_{i_0-1}(x+h)}{U_{i_0-2}(x+h)} \frac{W(f_1, f_2)}{g(x+h)} = S(x) \frac{W(f_1, f_2)}{g(x+h)}. \end{aligned}$$

Since the space spanned by $u_1, \dots, u_{i_0} \in V$ is acceptable, the polynomial $g(x+h) = W(u_1, \dots, u_{i_0-2})(x+h)/U_{i_0-2}(x+h)$ and $S(x)$ are relatively prime. Therefore the Wronskian $W(u_1, \dots, u_{i_0})$ is divisible by $S(x)$. $\square$

4.4. Frames of the fundamental spaces. Let $\mathbf{y} = (y_1, \dots, y_N)$ represent an h -critical point associated to the initial data $\mathbf{z}, \mathbf{\Lambda}, \mathbf{b}^{(i)}$. Let P be a population of h -critical points originated at $\mathbf{y}$.

Proposition 4.10. *The fundamental space V_P of the population P has no base points. The polynomials $T_1(x), \dots, T_N(x)$ given by (3.4) form a frame of V_P .*

Proof. We construct polynomials $u_1, \dots, u_{N+1} \in V_P$ as in the proof of Proposition 4.7. We have

$$\begin{aligned} W(u_1, \dots, u_i) &= y_i(x) \prod_{j=1}^{i-1} \prod_{r=1}^{i-j} T_j(x + (r-1)h), \\ W(u_1, \dots, u_{i-1}, u_{i+1}) &= \tilde{y}_i(x) \prod_{j=1}^{i-1} \prod_{r=1}^{i-j} T_j(x + (r-1)h), \end{aligned}$$

where $(y_1, \dots, \tilde{y}_i, \dots, y_N)$ is a descendant of $\mathbf{y}$ in the i direction.

In particular $W(u_1, \dots, u_i)$ is divisible by $\prod_{j=1}^{i-1} \prod_{r=1}^{i-j} T_j(x + (r-1)h)$. Therefore Wronskians of all i dimensional planes are divisible by this polynomial.

Moreover, we have $W(y_i, \tilde{y}_i) = T_i(x)y_{i-1}(x+h)y_{i+1}(x)$. Since y_i and $T_i(x)y_{i-1}(x+h)y_{i+1}(x)$ have no common roots, the polynomials $y_i(x)$ and $\tilde{y}_i(x)$ have no common roots. It follows that the greatest common divisor of i -dimensional Wronskians is $\prod_{j=1}^{i-1} \prod_{r=1}^{i-j} T_j(x + (r-1)h)$.

The absence of base points for V_P follows from the case of $i = 1$. $\square$

The converse statement is also true.

Proposition 4.11. *Let V be a space of polynomials of dimension $N + 1$ without base points and with the frame T_i . Let $\mathbf{z}, \mathbf{\Lambda}, \mathbf{b}^{(i)}$ be the initial data related to polynomials T_i by (3.4). Then V is the fundamental space of a population of h -critical points associated to the initial data $\mathbf{z}, \mathbf{\Lambda}, \mathbf{b}^{(i)}$.*

Proof. We postpone the proof until after Section 4.6, where the generating morphism β is described. Then the proof is similar to the proof of Proposition 5.17 in [MV1] combined with Lemma 5.20 in [MV1]. $\square$

From Propositions 4.10 and 4.11 we obtain the following theorem.

Theorem 4.12. *There is a bijective correspondence between populations of h -critical points associated to a given initial data and the spaces of polynomials with framing T_i , where T_i are related to the initial data by (3.4). $\square$*

4.5. Schubert calculus. The problem of counting the number of populations for a special choice of relative shifts $\mathbf{b}^{(i)}$ can be approached via Schubert calculus.

Let $\mathcal{V}$ be a complex vector space of dimension $d + 1$ and

$$\mathcal{F} = \{0 \subset F_1 \subset F_2 \subset \dots \subset F_{d+1} = \mathcal{V}\}, \quad \dim F_i = i,$$

a full flag in $\mathcal{V}$. Let $Gr(N + 1, \mathcal{V})$ be the Grassmanian of all $(N + 1)$ -dimensional subspaces in $\mathcal{V}$.

Let $\mathbf{a} = (a_1, \dots, a_{N+1})$, $d - N \geq a_1 \geq a_2 \geq \dots \geq a_{N+1} \geq 0$, be a non-increasing sequence of non-negative integers. Define the *Schubert cell* $G_{\mathbf{a}}^0(\mathcal{F})$ associated to the flag $\mathcal{F}$ and sequence $\mathbf{a}$ as the set

$$\begin{aligned} \{V \in Gr(N + 1, \mathcal{V}) \mid & \dim(V \cap F_{d-N+i-a_i}) = i, \\ & \dim(V \cap F_{d-N+i-a_i-1}) = i - 1, \text{ for } i = 1, \dots, N + 1\}. \end{aligned}$$

The closure $G_{\mathbf{a}}(\mathcal{F})$ of the Schubert cell is called *the Schubert cycle*. For a fixed flag $\mathcal{F}$, the Schubert cells form a cell decomposition of the Grassmanian. The codimension of $G_{\mathbf{a}}^0(\mathcal{F}(z))$ in the Grassmanian is $|\mathbf{a}| = a_1 + \dots + a_{N+1}$.

Let $\mathcal{V} = \mathbb{C}_d[x]$ be the space of polynomials of degree not greater than d , $\dim \mathcal{V} = d + 1$. For any $z \in \mathbb{C} \cup \infty$, define a full flag in $\mathbb{C}_d[x]$,

$$\mathcal{F}(z) = \{0 \subset F_1(z) \subset F_2(z) \subset \dots \subset F_{d+1}(z) = \mathcal{V}\}.$$

For $z \in \mathbb{C}$ and any i , we set $F_i(z)$ to be the subspace of all polynomials divisible by $\prod_{j=1}^{d+1-i} (x - z - (j-1)h)$. For any i , we set $F_i(\infty)$ to be the subspace of all polynomials of degree less than i .

Let $V \in Gr(N+1, \mathbb{C}_d[x])$. For any $z \in \mathbb{C} \cup \infty$, let $\mathbf{a}(z)$ be such a unique sequence that V belongs to the cell $G_{\mathbf{a}(z)}^0(\mathcal{F}(z))$. We say that a point $z \in \mathbb{C} \cup \infty$ is an h -ramification point for V , if $\mathbf{a}(z) \neq (0, \dots, 0)$. We call $\mathbf{a}(z)$ the *ramification condition of V at z* .

If $u_1, \dots, u_{N+1}$ is a basis of V then the ramification points are zeroes of the discrete Wronskian $W(x) = W(u_1, \dots, u_{N+1})(x)$. Indeed, given $z \in \mathbb{C}$, without loss of generality we can assume that $u_i \in \mathcal{F}_{d+2-i-a_i(z)}(z)$ and therefore the matrix $\{u_i(x + (j-1)h)\}_{i,j=1}^{N+1}$ is upper triangular. Moreover, this matrix has a zero diagonal element if and only if $|a| > 0$.

Fix h -ramification conditions at $z_1, \dots, z_n, \infty$ so that

$$\sum_{s=1}^n |\mathbf{a}(z_s)| + |\mathbf{a}(\infty)| = \dim Gr(N+1, \mathbb{C}_d[x]) .$$

Counting Problem. *Compute the number of spaces of polynomials of dimension $N+1$ with these ramification properties.*

In other words we ask to count the number of points in the intersection of Schubert cycles. The intersection index of the Schubert cycles is well known.

Namely, for a non-increasing sequence $\mathbf{a} = (a_1, \dots, a_{N+1})$, $a_1 \geq a_2 \geq \dots \geq a_{N+1} \geq 0$, of non-negative integers, denote $\tilde{L}_{\mathbf{a}}$ the finite dimensional irreducible gl_{N+1} -module with highest weight $\mathbf{a}$. Let $L_{\mathbf{a}}$ be the sl_{N+1} module obtained by restriction of $\tilde{L}_{\mathbf{a}}$.

Theorem 4.13. ([F]) *The intersection index of Schubert cycles $G_{\mathbf{a}(z_s)}(F(z_s))$, $s = 1, \dots, n$, and $G_{\mathbf{a}(\infty)}(F(\infty))$ equals the multiplicity of the trivial sl_{N+1} -module in the tensor product of sl_{N+1} -modules*

$$L_{\mathbf{a}(z_1)} \otimes \dots \otimes L_{\mathbf{a}(z_n)} \otimes L_{\mathbf{a}(\infty)} .$$

Conjecturally, for almost all $z_1, \dots, z_n$ the number of spaces V with such ramification conditions is equal to the above multiplicity.

Fix an initial data $\mathbf{z}, \mathbf{\Lambda}, \mathbf{b}^{(i)}$. Until the end of this section we assume that $z_s - z_r \notin h\mathbb{Z}$ for all $s \neq r$ and

$$b_s^{(j)} = - \sum_{i=1}^j \Lambda_s^{(i)} . \quad (4.4)$$

Fix an sl_N weight and write it in the form $w \cdot \Lambda_{\infty}$ where Λ_{∞} is dominant integral and w is an element of Weyl group.

Let $\mathbf{y}$ be an h -critical point associated to the initial data and the weight at infinity $w \cdot \Lambda_{\infty}$. Let P be the population of h -critical points originated at $\mathbf{y}$ and let V_P be the corresponding fundamental space. Let d be large enough, so that $V_P \subset \mathbb{C}_d[x]$.

Theorem 4.14. *The points $z_1, \dots, z_n$ and ∞ are ramification points of V_P . The ramification condition at z_s is $a_i(z_s) = \sum_{j=1}^{N+1-i} \Lambda_s^{(j)}$. The ramification condition at ∞ is $a_i(\infty) = d - N - l_1 - \sum_{j=1}^{i-1} \Lambda_\infty^{(j)}$, where l_1 is defined from Λ and Λ_∞ via (3.1).*

Proof. Follows from Lemma 4.8, cf. proof of Lemmas 5.8, 5.10 in [MV1]. $\square$

For an integral dominant sl_{N+1} weight Λ , we denote L_Λ the irreducible sl_{N+1} module with highest weight Λ .

Corollary 4.15. *The number of (discretely lying) populations associated to initial data $\mathbf{z}_i, \Lambda, \mathbf{b}^{(i)}$ such that (4.4) holds, is not greater than the multiplicity of L_{Λ_∞} in the tensor product $L_{\Lambda_1} \otimes \dots \otimes L_{\Lambda_n}$.*

Proof. The corollary holds since the number of isolated points of the intersection of the corresponding Schubert cycles is not greater than the intersection index of the cycles. $\square$

4.6. Generating morphism. We identify a population with the variety of full flags in the fundamental space.

Let $V = V_P$ be the fundamental space of a population P . By Proposition 4.10, the polynomials T_i defined in (3.4) form a frame of V_P .

Let $FL(V)$ be the variety of all full flags

$$\mathcal{F} = \{0 \subset F_1 \subset F_2 \subset \dots \subset F_{N+1} = V\}$$

in V . For any $\mathcal{F} \in FL(V)$ define an N -tuple of polynomials $\mathbf{y}^\mathcal{F} = (y_1^\mathcal{F}, \dots, y_N^\mathcal{F})$ as follows. Let $u_1, \dots, u_{N+1}$ be a basis in V such that for any i the polynomials $u_1, \dots, u_i$ form a basis in F_i . We say that this basis is *adjusted* to the flag $\mathcal{F}$ and the flag $\mathcal{F}$ is *generated* by the basis $u_1, \dots, u_{N+1}$.

Define the polynomials

$$y_i^\mathcal{F}(x) = \frac{W(u_1, \dots, u_i)(x)}{\prod_{j=1}^{i-1} T_1(x + (j-1)h) \prod_{j=1}^{i-2} T_2(x + (j-1)h) \dots T_{i-1}(x)}. \quad (4.5)$$

The correspondence $\mathcal{F} \mapsto \mathbf{y}^\mathcal{F}$ gives a morphism

$$\beta : FL(V) \rightarrow \mathbf{P}(\mathbb{C}[x])^N, \quad (4.6)$$

called *the generating morphism of V* .

Theorem 4.16. *Let P be a population of h -critical points with the fundamental space V . Then the generating morphism defines an isomorphism of $FL(V)$ and the population $P \subset \mathbf{P}(\mathbb{C}[x])^N$.*

Proof. Proof is the same as the proof of the first part of Theorem 5.12 in [MV1]. $\square$

Remark. Recall that V is the $(N+1)$ -dimensional complex vector space of polynomials which is contained in the kernel of the fundamental operator D_P and D_P is a

linear difference operator of order $N + 1$. Therefore the full flags in V also label the decompositions of D_P to $N + 1$ linear factors of the form $(\partial - f(x))$, where $f(x)$ is a rational function of x . Thus, h -critical points are in bijective correspondence with such factorizations of the fundamental operator D_P .

Fix a flag $\mathcal{F}^0 \in FL(V)$. For any $\mathcal{F} \in FL(V)$ define a permutation $w(\mathcal{F})$ in the symmetric group S^{N+1} as follows. Define $w_1(\mathcal{F})$ as the minimum of i such that $F_1 \subset F_i^0$. Fix a basis vector $u_1 \in F_1$. Define $w_2(\mathcal{F})$ as the minimum of i such that there is a basis in F_2 of the form u_1, u_2 with $u_2 \in F_i^0$. Assume that $w_1(\mathcal{F}), \dots, w_j(\mathcal{F})$ and $u_1, \dots, u_j$ are determined. Define $w_{j+1}(\mathcal{F})$ as the minimum of i such that there is a basis in F_{j+1} of the form $u_1, \dots, u_j, u_{j+1}$ with $u_{j+1} \in F_i^0$. As a result of this procedure we define $w(\mathcal{F}) = (w_1(\mathcal{F}), \dots, w_{N+1}(\mathcal{F})) \in S^{N+1}$ and a basis $u_1, \dots, u_{N+1}$ which generates $\mathcal{F}$ and such that $u_i \in F_{w_i(\mathcal{F})}^0$.

For $w \in S^{N+1}$, define

$$G_w^{\mathcal{F}^0} = \{\mathcal{F} \in FL(V), w(\mathcal{F}) = w\}.$$

The algebraic variety $G_w^{\mathcal{F}^0}$ is called *the Bruhat cell* associated with $\mathcal{F}^0$ and $w \in S^{N+1}$. The set of all Bruhat cells form a cell decomposition of $FL(V)$:

$$FL(V) = \sqcup_{w \in S^{N+1}} G_w^{\mathcal{F}^0}.$$

Recall that the symmetric group S^{N+1} is identified with the sl_{N+1} Weyl group in such a way that the simple transposition $(i, i + 1)$ corresponds to the simple reflection with respect to α_i .

Now we are ready to describe the set of $(N + 1)$ -tuples in a population of a fixed degree. Let $\mathbf{y}$ represent an h -critical point associated with the initial data and the weight at infinity $w \cdot \Lambda_\infty$, where Λ_∞ is dominant integral. Let $P = P(\mathbf{y})$ be the corresponding population and $\beta : FL(V) \rightarrow P$ the generating isomorphism.

Theorem 4.17. *The set of N -tuples $\bar{\mathbf{y}}$ in $P(\mathbf{y})$ associated with a weight at infinity $w \cdot \Lambda_\infty$, where Λ_∞ is integral dominant, coincides with the image of the Bruhat cell $\beta(G_w^{\mathcal{F}(\infty)})$.*

Proof. The theorem is proved similar to Corollary 5.23 in [MV1]. □

In particular, each population contains at most one tuple $\mathbf{y}$ which represents an h -critical point associated to a integral dominant weight at infinity. Thus the number of critical points associated to an initial data and an integral dominant weight is bounded from above by the number of the populations. We conjecture that for generic values of z_i this bound is exact.

5. h -SELF DUAL VECTOR SPACES OF POLYNOMIALS

5.1. Dual spaces. Let V be a space of polynomials of dimension $N + 1$ with frame $T_1(x), \dots, T_N(x)$. For $u_1, \dots, u_i \in V$, the polynomial

$$W^\dagger(u_1, \dots, u_i)(x) := \frac{W(u_1, \dots, u_i)(x)}{\prod_{j=1}^{i-1} T_1(x + (j-1)h) \prod_{j=1}^{i-2} T_2(x + (j-1)h) \dots T_{i-1}(x)}$$

is called *the divided Wronskian*.

Note that if polynomials $u_1, \dots, u_{N+1}$ form a basis of V then the divided Wronskian $W^\dagger(u_1, \dots, u_{N+1})$ is a non-zero constant.

Let $V^\dagger$ be the set of polynomials $W^\dagger(u_1, \dots, u_N)$, where $u_i \in V$. Clearly $V^\dagger$ is a vector space of dimension $N + 1$. We call $V^\dagger$ the *h -dual space of V* .

We have a non-degenerate pairing

$$V \otimes V^\dagger \rightarrow \mathbb{C}, \quad u \otimes W^\dagger(u_1, \dots, u_N) \mapsto W^\dagger(u, u_1, \dots, u_N)(x).$$

For $i = 1, \dots, N$, set

$$T_i^\dagger(x) = T_{N+1-i}(x + (i-1)h). \quad (5.1)$$

The following two lemmas are obtained from Wronskian identities in Lemmas 9.5 and 9.3.

Lemma 5.1. *The polynomials $T_1^\dagger(x), \dots, T_N^\dagger(x)$ form a frame of $V^\dagger$.* $\square$

For a space of polynomials V and $a \in \mathbb{C}$, we denote $V(x+a)$ the space of polynomials spanned by $u(x+a)$, $u(x) \in V$.

Lemma 5.2. *We have $V^{\dagger\dagger} = V(x + (N-1)h)$.* $\square$

5.2. h -selfdual spaces and canonical bilinear form. We say that V is *h -selfdual*, if $V^\dagger = V(x - (N-1)h/2)$.

A space V is *h -selfdual* if and only if $V^\dagger$ is *h -selfdual*.

If V is *h -selfdual*, then

$$T_i(x) = T_{N+1-i}(x + (i-1)h - (N-1)h/2). \quad (5.2)$$

In particular, if T_i are of the form (3.4), then

$$\Lambda_s^{(i)} = \Lambda_s^{(N+1-i)}, \quad b_s^{(i)} = b_s^{(N+1-i)} + (i-1)h - (N-1)h/2.$$

For instance, the space of polynomials of degree not greater than N is *h -selfdual*. In this case all polynomials T_i are equal to 1.

Let V be *h -selfdual*. Define a non-degenerate pairing $(\ , \) : V \otimes V \rightarrow \mathbb{C}$. If $u, v \in V$, then we write $v(x) = W^\dagger(u_1, \dots, u_N)(x - (N-1)h/2)$ with $u_i \in V$ and set $(u, v) = W^\dagger(u, u_1, \dots, u_N)$. This pairing is called the *canonical bilinear form*.

A basis $u_1, \dots, u_{N+1}$ in a space of polynomials V is called a *Witt basis* if for $i = 1, \dots, N+1$ we have

$$u_i(x) = W^\dagger(u_1, \dots, \widehat{u}_{N+2-i}, \dots, u_{N+1})(x - (N-1)h/2). \quad (5.3)$$

Clearly, if V has a Witt basis, then V is h -selfdual.

Theorem 5.3. *Let V be h -selfdual. Then V has a Witt basis. In particular, the form $(\ , \) : V \otimes V \rightarrow \mathbb{C}$ is symmetric if the dimension of V is odd and skew-symmetric if the dimension of V is even.*

Proof. The theorem is proved similarly to Theorem 6.4 in [MV1]. $\square$

5.3. Isotropic flags. Let $u_1, \dots, u_{N+1}$ be a basis in a vector space V of polynomials. Denote $W_i(x) = W^\dagger(u_1, \dots, \widehat{u_i}, \dots, u_{N+1})(x - (N - 1)h/2)$.

Lemma 5.4. *If for $i = 1, \dots, N$*

$$W^\dagger(u_1, \dots, u_i)(x) = c_i W^\dagger(u_1, \dots, u_{N+1-i})(x + (i - 1)h - \frac{N - 1}{2}h), \quad (5.4)$$

where $c_1, \dots, c_N$ are some non-zero complex numbers, then for every i , the polynomial $u_i(x)$ is a linear combination of $W_{N+1}(x), W_N(x), \dots, W_{N+2-i}(x)$.

Proof. The proof of the lemma is similar to the proof of Theorem 6.8 in [MV1]. $\square$

Corollary 5.5. *If V has a basis satisfying (5.4), then V is h -selfdual.*

Let V be an h -selfdual space of polynomials. For a subspace $U \subset V$ denote $U^\perp$ its orthogonal complement. A full flag $\mathcal{F} = \{0 \subset F_1 \subset \dots \subset F_{N+1} = V\}$ is called *isotropic* if $F_i^\perp = F_{N+1-i}$ for $i = 1, \dots, N$.

Proposition 5.6. *Let $\mathcal{F}$ be a flag in an h -selfdual space of polynomials V and $u_1, \dots, u_{N+1}$ a basis in V adjusted to $\mathcal{F}$. Then $\mathcal{F}$ is isotropic if and only if (5.4) holds.*

Proof. If $F_i^\perp = F_{N+1-i}$ then we have two bases in F_{N+1-i} : the basis $u_1, \dots, u_{N+1-i}$ and the basis $W_{N+1}, W_N, \dots, W_{i+1}$. Hence

$$\begin{aligned} W^\dagger(u_1, \dots, u_{N+1-i})(x) &= \text{const} W^\dagger(W_{N+1}, W_N, \dots, W_{i+1})(x) = \\ &= \text{const} W^\dagger(u_1, \dots, u_i)(x + (N - i)h - (N - 1)h/2). \end{aligned}$$

The only if part of the lemma is proved.

Now, let (5.4) hold. We prove that the spaces spanned by $u_1(x), \dots, u_i(x)$ and by $W_{N+1}(x - (N - 1)h/2), \dots, W_{N-i}(x - (N - 1 + 2)h/2)$ are the same by induction on i . For $i = 1$ it is just equation (5.4). Assume that the statement is proved for $i = 1, \dots, i_0 - 1$. Then

$$\begin{aligned} W^\dagger(u_1, \dots, u_{i_0})(x) &= \text{const} W^\dagger(u_1, \dots, u_{N+1-i_0})(x + (i_0 - 1)h - (N - 1)h/2) = \\ &= \text{const} W^\dagger(W_{N+1}, W_N, \dots, W_{N-i_0+2})(x - (N - 1)h/2) = \\ &= \text{const} W^\dagger(u_1(x), \dots, u_{i_0-1}(x), W_{N-i_0+2}(x - (N - 1)h/2)), \end{aligned}$$

and the step of induction follows. $\square$

6. THE CASE OF B_N

Consider the root system of type B_N corresponding to the Lie algebra so_{2N+1} . Let $\alpha_1, \dots, \alpha_{N-1}$ be long simple roots and α_N the short one. We have

$$(\alpha_N, \alpha_N) = 2, \quad (\alpha_i, \alpha_i) = 4, \quad (\alpha_i, \alpha_{i+1}) = -2,$$

for $i = 1, \dots, N-1$, and all other scalar products are zero.

6.1. Definition of critical points of B_N type. We fix ramification points $\mathbf{z} = (z_1, \dots, z_n) \in \mathbb{C}^n$, non-zero dominant integral B_N weights $\mathbf{\Lambda} = (\Lambda_1, \dots, \Lambda_n)$ and relative shifts $\mathbf{b}^{(i)} = (b_1^{(i)}, \dots, b_n^{(i)}) \in \mathbb{C}^n$, $i = 1, \dots, N$. We call this data a B_N initial data. We also set $\Lambda_s^{(i)} = (\Lambda_s, \alpha_i^\vee)$.

Given $\mathbf{l} = (l_1, \dots, l_N) \in \mathbb{Z}_{\geq 0}^N$, we define the B_N weight at infinity Λ_∞ by the formula (3.1).

For a B_N initial data $\mathbf{z}, \mathbf{\Lambda}, \mathbf{b}^{(i)}$ and a weight at infinity Λ_∞ , we define an sl_{2N} initial data which consists of ramification points z_i , non-zero dominant integral sl_{2N} weights Λ_i^A , relative shifts $\mathbf{b}^{(i),A}$ and we also define an sl_{2N} weight at infinity Λ_∞^A . Namely, given a B_N weight Λ , the sl_{2N} weight Λ^A is defined by

$$(\Lambda^A, \alpha_i^A) = (\Lambda^A, \alpha_{2N-i}^A) = (\Lambda, \alpha_i^\vee),$$

where $i = 1, \dots, N$ and α_i^A are roots of sl_{2N} . The shifts $\mathbf{b}^{(i),A}$ are defined by

$$b_s^{(i),A} = b_s^{(2N-i),A} + (i - N)h = b_s^{(i)}.$$

Given a set of complex numbers $t_j^{(i)}$, $i = 1, \dots, N$, $j = 1, \dots, l_i$, we represent it by the N -tuple of polynomials $\mathbf{y} = (y_1, \dots, y_N)$, where $y_i(x) = \prod_{j=1}^{l_i} (x - t_j^{(i)})$. We define the corresponding $(2N-1)$ -tuple $\mathbf{y}^A$ by the formula

$$y_i^A(x) = y_i(x), \quad y_N^A(x) = y_N(x), \quad y_{i+N}^A(x) = y_{N-i}(x + ih), \quad (6.1)$$

where $i = 1, \dots, N-1$.

We propose the following definition of the h -critical points of type B_N , cf. Lemma 7.1 in [MV1]. The set of complex numbers $t_j^{(i)}$, $i = 1, \dots, N$, $j = 1, \dots, l_i$, is called an h -critical point of B_N type associated to the initial data $\mathbf{z}, \mathbf{\Lambda}, \mathbf{b}^{(i)}$ and the weight at infinity Λ_∞ if the corresponding $(2N-1)$ -tuple $\mathbf{y}^A$ represents an sl_{2N} h -critical point associated to the initial data $\mathbf{z}, \mathbf{\Lambda}^A, \mathbf{b}^{(i),A}$ and the weight at infinity Λ_∞^A .

In this case we say that the N -tuple $\mathbf{y}$ represents an h -critical point of the B_N type.

Equivalently, the set of numbers $t_j^{(i)}$, $i = 1, \dots, N$, $j = 1, \dots, l_i$, is an h -critical point of B_N type if it satisfies the following system of algebraic equations:

$$\prod_{s=1}^n \frac{t_j^{(i)} - z_s + b_s^{(i)}h + \Lambda_s^{(i)}h}{t_j^{(i)} - z_s + b_s^{(i)}h} \prod_{k=1}^{l_{i-1}} \frac{t_j^{(i)} - t_k^{(i-1)} + h}{t_j^{(i)} - t_k^{(i-1)}} \prod_{k \neq j} \frac{t_j^{(i)} - t_k^{(i)} - h}{t_j^{(i)} - t_k^{(i)} + h} \times$$

$$\times \prod_{k=1}^{l_{i+1}} \frac{t_j^{(i)} - t_k^{(i+1)}}{t_j^{(i)} - t_k^{(i+1)} - h} = 1,$$

$$\prod_{s=1}^n \frac{t_j^{(N)} - z_s + b_s^{(N)}h + \Lambda_s^{(N)}h}{t_j^{(N)} - z_s + b_s^{(N)}h} \prod_{k=1}^{l_{N-1}} \left(\frac{t_j^{(N)} - t_k^{(N-1)} + h}{t_j^{(N)} - t_k^{(N-1)}} \right)^2 \prod_{k \neq j} \frac{t_j^{(N)} - t_k^{(N)} - h}{t_j^{(N)} - t_k^{(N)} + h} = 1,$$

where $i = 1, \dots, N-1$.

Note that in the quasiclassical limit $h \rightarrow 0$, this system becomes system (2.2) of [MV1] specialized to the case of B_N .

6.2. Reproduction procedure of B_N type. We describe the concepts of reproduction and population. All of that follows from the definitions in the case of sl_{2N} .

An N -tuple $\mathbf{y}$ is called fertile in B_N sense if there exist polynomials $\tilde{y}_i$, $i = 1, \dots, N-1$ and $\tilde{y}_N$ such that

$$\begin{aligned} W(y_i, \tilde{y}_i)(x) &= y_{i+1}(x)y_{i-1}(x+h)T_i(x), \\ W(y_N, \tilde{y}_N)(x) &= y_{N-1}^2(x+h)T_N(x). \end{aligned}$$

The N -tuple $\mathbf{y}^{(i)} = (y_1, \dots, \tilde{y}_i, \dots, y_N)$ is called an immediate descendant of $\mathbf{y}$ in the direction i . Immediate descendants $\mathbf{y}^{(i)}$ are also fertile in B_N sense.

From Lemma 3.3 we obtain that an N -tuple $\mathbf{y}$ represents a critical point if and only if it is fertile and $\mathbf{y}^A$ is generic. In this case if $(\mathbf{y}^{(i)})^A$ is generic, then $\mathbf{y}^{(i)}$ also represents an h -critical point of type B_N .

The B_N population originated at a critical point $\mathbf{y}$ is the minimal set $P(\mathbf{y})$ of fertile N -tuples such that $\mathbf{y} \in P(\mathbf{y})$ and if $\bar{\mathbf{y}} \in P(\mathbf{y})$ then the immediate descendants $\bar{\mathbf{y}}^{(i)} \in P(\mathbf{y})$.

Obviously, the B_N population is contained in the corresponding sl_{2N} population: if $\bar{\mathbf{y}} \in P(\mathbf{y})$ then $\bar{\mathbf{y}}^A \in P_A(\mathbf{y}^A)$, where we denote $P_A(\mathbf{y}^A)$ the sl_{2N} population of critical points originated at $\mathbf{y}^A$. The fundamental space of the sl_{2N} population $P_A(\mathbf{y}^A)$ contains a flag with the property (5.3) and therefore is h -selfdual. We call this space the *fundamental space of the B_N population $P(\mathbf{y})$* .

The corresponding fundamental operator is the difference operator of order $2N$ obtained from (4.2) by substituting (6.1) and (5.2):

$$\prod_i^{N-1} \left(\partial - \frac{y_{N-i}(x + (i+1)h)}{y_{N-i}(x + ih)} \frac{y_{N+1-i}(x + (i-1)h)}{y_{N+1-i}(x + ih)} T(x + ih) \prod_{s=1}^{i-1} \frac{T_{N-s}(x + ih)}{T_{N-s}(x + (i-1)h)} \right) \times \\ \prod_i^{1 \rightarrow N} \left(\partial - \frac{y_{N+1-i}(x + h)}{y_{N+1-i}(x)} \frac{y_{N-i}(x)}{y_{N-i}(x + h)} \prod_{s=1}^{N-i} \frac{T_s(x + (N-i-s+1)h)}{T_s(x + (N-i-s)h)} \right),$$

where $T(x) = \prod_{s=1}^N T_s(x + (N-s)h) / T_s(x + (N-s-1)h)$.

Clearly, two populations of h -critical points of type B_N either do not intersect or coincide.

6.3. A B_N population and the C_N flag variety. Let $\mathbf{y}$ be an h -critical point of B_N type and let V be the fundamental space of the population $P_A(\mathbf{y}^A)$. Then V is an h -selfdual space of dimension $2N$ and therefore has a non-degenerate skew-symmetric canonical bilinear form.

The special symplectic Lie algebra of V consists of all traceless endomorphisms x of V such that $(xv, v') + (v, xv') = 0$ for all $v, v' \in V$. Let $\mathbf{u} = (u_1, \dots, u_{2N})$ be a Witt basis in V . We have $(u_i, u_{2N+1-i}) = (-1)^{i+1}$, $i = 1, \dots, N$, and $(u_i, u_j) = 0$ if $i + j \neq 2N + 1$. This choice of basis identifies the special symplectic Lie algebra with a Lie subalgebra of sl_{2N} , which is denoted sp_{2N} . The Lie algebra sp_{2N} has the root system of type C_N .

Denote $E_{i,j}$ the matrix with zero entries except 1 at the intersection of the i -th row and the j -th column. The lower triangular part of sp_{2N} is spanned by matrices $E_{i,i+1} + E_{2N-i, 2N+1-i}$ for $i = 1, \dots, N-1$ and $E_{N, N+1}$. Denote these matrices $X_1, \dots, X_N$, respectively.

Now we describe the action of one-parametric subgroups in the special symplectic group in the direction of X_i on the basis $\mathbf{u} = (u_1, \dots, u_{2N})$. We have

$$e^{cX_i} \mathbf{u} = \\ = (u_1, \dots, u_{i-1}, u_i + cu_{i+1}, u_{i+1}, \dots, u_{2N-i}, u_{2N+1-i} + cu_{2N+2-i}, u_{2N+2-i}, \dots, u_{2N}), \\ e^{cX_i} \mathbf{u} = (u_1, \dots, u_{N-1}, u_N + cu_{N+1}, u_{N+1}, \dots, u_{2N}),$$

where $i = 1, \dots, N-1$.

We also set

$$e^{\infty X_i} \mathbf{u} = (u_1, \dots, u_{i-1}, u_{i+1}, u_i, \dots, u_{2N-i}, u_{2N+2-i}, u_{2N+1-i}, \dots, u_{2N}), \\ e^{\infty X_i} \mathbf{u} = (u_1, \dots, u_{N-1}, u_{N+1}, u_N, \dots, u_{2N}).$$

Note $e^{cX_i} \mathbf{u}$ is a Witt basis for all i, c and the corresponding flag is isotropic.

Given a basis $\mathbf{u} = (u_1, \dots, u_{2N})$ of V we denote $\mathcal{F}(\mathbf{u})$ the full flag generated by this basis. Let $FL(V)$ be the variety of all full flags of V .

Denote $FL^\perp(V) \subset FL(V)$ the subvariety of all isotropic flags and define the B_N generating morphism $\beta : FL^\perp(V) \rightarrow \mathbf{P}(\mathbb{C}[x])^N$, $\mathcal{F}(\mathbf{u}) \mapsto (y_1^\mathcal{F}, \dots, y_N^\mathcal{F})$, where $y_i^\mathcal{F} = W^\dagger(u_1, \dots, u_i)$.

Lemma 6.1. *If $F(\mathbf{u})$ is an isotropic flag then $\beta(\mathcal{F}(\mathbf{u}))$ is fertile. Moreover the set of all immediate descendants of $\beta(\mathcal{F}(\mathbf{u}))$ in the direction i coincides with the set $\beta(\mathcal{F}(e^{cX_i}\mathbf{u}))$, $c \in \bar{\mathbb{C}}$.*

Proof. Follows from the Wronskian identities

$$\begin{aligned} W(y_i^\mathcal{F}, W^\dagger(u_1, \dots, u_{i-1}, u_{i+1}))(x) &= T_i(x) y_{i-1}^\mathcal{F}(x+h) y_{i+1}^\mathcal{F}(x), \\ W(y_N^\mathcal{F}, W^\dagger(u_1, \dots, u_{N-1}, u_{N+1}))(x) &= T_N(x) (y_{N-1}^\mathcal{F}(x+h))^2, \end{aligned}$$

where $i = 1, \dots, N-1$. □

Theorem 6.2. *The generating morphism β of type B_N gives rise to the isomorphism $FL^\perp(V) \rightarrow P(\mathbf{y}) \subset \mathbf{P}(\mathbb{C}[x])^N$.*

Proof. The generating morphism β is an isomorphism of $FL^\perp(V)$ to the image by Theorem 4.16. Clearly, the image of β contains the population $P(\mathbf{y})$. The image of β coincides with $P(\mathbf{y})$ by Lemma 6.1. □

Note that $FL^\perp$ is isomorphic to the flag variety of the special symplectic group which corresponds to the C_N root system. This flag variety has a Bruhat cell decomposition $FL^\perp(V) = \sqcup_{w \in \mathcal{W}} G_w^{\mathcal{F}(\infty), C}$, where $\mathcal{W}$ is the C_N Weyl group. Note that the Weyl groups of B_N and C_N are canonically identified. See for details, e.g. [MV1], section 7.3.

As before, the set of all N -tuples in a B_N population of the same degree is the Bruhat cell in the C_N flag variety.

Theorem 6.3. *The set of N -tuples $\bar{\mathbf{y}}$ in $P(\mathbf{y})$ associated with a weight at infinity $w \cdot \Lambda_\infty$, where Λ_∞ is integral dominant, coincides with the image of the Bruhat cell $\beta(G_w^{\mathcal{F}(\infty), C})$.*

Proof. Completely parallel to the proof of Corollary 7.12 in [MV1]. □

7. THE CASE OF C_N

Consider the root system of type C_N corresponding to the Lie algebra sp_{2N} . Let $\alpha_1, \dots, \alpha_{N-1}$ be short simple roots and α_N the long one. We have

$$(\alpha_N, \alpha_N) = 4, \quad (\alpha_i, \alpha_i) = 2, \quad (\alpha_{N-1}, \alpha_N) = -2 \quad (\alpha_{i-1}, \alpha_i) = -1,$$

for $i = 1, \dots, N-1$ and all other scalar products are zero.

7.1. Definition of critical points of C_N type. We fix ramification points $\mathbf{z} = (z_1, \dots, z_n) \in \mathbb{C}$, dominant integral C_N weights $\mathbf{\Lambda} = (\Lambda_1, \dots, \Lambda_n)$ and relative shifts $\mathbf{b}^{(i)} = (b_1^{(i)}, \dots, b_n^{(i)}) \in \mathbb{C}^n$, $i = 1, \dots, N$. We call this data a C_N initial data. We also set $\Lambda_s^{(i)} = (\Lambda_s, \alpha_i^\vee)$.

Given $\mathbf{l} = (l_1, \dots, l_N) \in \mathbb{Z}_{\geq 0}^N$, we define the C_N weight at infinity Λ_∞ by the formula (3.1).

For a C_N initial data $\mathbf{z}, \mathbf{\Lambda}, \mathbf{b}^{(i)}$ and a weight at infinity Λ_∞ , we define an sl_{2N+1} initial data which consists of ramification points z_i , dominant integral Λ_i^A , relative shifts $\mathbf{b}^{(i),A}$ and we also define an sl_{2N+1} weight at infinity Λ_∞^A . Namely, given a C_N weight Λ , the sl_{2N+1} weight Λ^A is defined by

$$(\Lambda^A, \alpha_i^A) = (\Lambda^A, \alpha_{2N+1-i}^A) = (\Lambda, \alpha_i^\vee),$$

where $i = 1, \dots, N$ and α_i^A are roots of sl_{2N+1} . The shifts $\mathbf{b}^{(i),A}$ are defined by

$$b_s^{(i),A} = b_s^{(2N+1-i),A} + (i - N + 1/2)h = b_s^{(i)}.$$

Given a set of complex numbers $t_j^{(i)}, t_k^{(N)}$ where $i = 1, \dots, N-1$, $j = 1, \dots, l_i$, $k = 1, \dots, 2l_N$, we represent it by the N -tuple of polynomials $\mathbf{y} = (y_1, \dots, y_N)$, where $y_i(x) = \prod_{j=1}^{l_i} (x - t_j^{(i)})$, $y_N = \prod_{j=1}^{2l_N} (x - t_j^{(N)})$. We define the corresponding $(2N)$ -tuple $\mathbf{y}^A$ by the formula

$$y_i^A(x) = y_i(x), \quad y_{i+N}^A(x) = y_{N+1-i}(x + (i - 1/2)h), \quad (7.1)$$

where $i = 1, \dots, N$.

We propose the following definition of the h -critical points of type C_N , cf. Section 7.2 in [MV1]. The set of complex numbers $t_j^{(i)}, t_k^{(N)}$, where $i = 1, \dots, N-1$, $j = 1, \dots, l_i$, $k = 1, \dots, 2l_N$, is called an h -critical point of C_N type associated to the initial data $\mathbf{z}, \mathbf{\Lambda}, \mathbf{b}^{(i)}$ and the weight at infinity Λ_∞ if the corresponding $(2N)$ -tuple $\mathbf{y}^A$ represents an sl_{2N+1} h -critical point associated to the initial data $\mathbf{z}, \mathbf{\Lambda}^A, \mathbf{b}^{(i),A}$ and the weight at infinity Λ_∞^A .

In this case we say that the N -tuple $\mathbf{y}$ represents an h -critical point of the C_N type.

Equivalently, the set of numbers $t_j^{(i)}, t_k^{(N)}$, where $i = 1, \dots, N-1$, $j = 1, \dots, l_i$, $k = 1, \dots, 2l_N$, is an h -critical point of C_N type if it satisfies the following system of

algebraic equations:

$$\begin{aligned}
& \prod_{s=1}^n \frac{t_j^{(i)} - z_s + b_s^{(i)}h + \Lambda_s^{(i)}h}{t_j^{(i)} - z_s + b_s^{(i)}h} \prod_k \frac{t_j^{(i)} - t_k^{(i-1)} + h}{t_j^{(i)} - t_k^{(i-1)}} \prod_{k \neq j} \frac{t_j^{(i)} - t_k^{(i)} - h}{t_j^{(i)} - t_k^{(i)} + h} \times \\
& \quad \times \prod_k \frac{t_j^{(i)} - t_k^{(i+1)}}{t_j^{(i)} - t_k^{(i+1)} - h} = 1, \\
& \prod_{s=1}^n \frac{t_j^{(N)} - z_s + b_s^{(N)}h + \Lambda_s^{(N)}h}{t_j^{(N)} - z_s + b_s^{(N)}h} \prod_{k=1}^{l_{N-1}} \frac{t_j^{(N)} - t_k^{(N-1)} + h}{t_j^{(N)} - t_k^{(N-1)}} \prod_{k \neq j} \frac{t_j^{(N)} - t_k^{(N)} - h}{t_j^{(N)} - t_k^{(N)} + h} \times \\
& \quad \times \prod_{k=1}^{2l_N} \frac{t_j^{(N)} - t_k^{(N)} + h/2}{t_j^{(N)} - t_k^{(N)} - h/2} = 1.
\end{aligned}$$

Note that in the quasiclassical limit $h \rightarrow 0$, this system becomes system (2.2) of [MV1] specialized to the case of C_N .

7.2. Reproduction procedure of C_1 type. Fix a polynomial $T(x)$. Suppose that we have a polynomial $y(x)$, such that $(y(x), y(x + h/2))$ is a critical point of sl_3 with weights and shifts given by $(T(x), T(x + h/2))$ via (3.4). Then there exists a polynomial $\tilde{y}$ such that

$$W(y, \tilde{y})(x) = y(x + h/2)T(x).$$

The fundamental space V of sl_3 population P_A originated at $(y(x), y(x + h/2))$ is 3-dimensional. It is spanned by $u_1 = y(x)$, $u_2 = \tilde{y}(x)$ and u_3 satisfying the identities

$$\begin{aligned}
W(u_1, u_2)(x) &= u_1(x + h/2)T(x), \\
W(u_1, u_3)(x) &= u_2(x + h/2)T(x), \\
W(u_2, u_3)(x) &= u_3(x + h/2)T(x).
\end{aligned} \tag{7.2}$$

These three equations constitute the fact that u_1, u_2, u_3 form a Witt basis of V .

Equations (7.2) imply the following lemma.

Lemma 7.1. *Let v be a polynomial in V . The pair $(v(x), v(x + h/2))$ is in the population P_A if and only if $v(x)$ is a scalar multiple of $u_1 + \alpha u_2 + \alpha^2 u_3/2$ for some $\alpha \in \mathbb{C}$ or of u_3 (i.e., $\alpha = \infty$). $\square$*

We say that the polynomials $v(x)$ described in the lemma and considered as elements of $\mathbf{P}(\mathbb{C}[x])$ form a C_1 population of h -critical points with weight $T(x)$. Note that unlike the case of sl_2 a C_1 population is not a linear space.

7.3. Reproduction procedure of C_N type. We describe the concepts of reproduction and population. All of that follows from the definitions in the cases of sl_{2N+1} and C_1 .

An N -tuple $\mathbf{y}$ is called fertile in C_N sense if there exist polynomials $\tilde{y}_i$, $i = 1, \dots, N-1$ and $\bar{y}_N$ such that

$$\begin{aligned} W(y_i, \tilde{y}_i)(x) &= y_{i+1}(x)y_{i-1}(x+h)T_i(x), \\ W(y_N, \bar{y}_N)(x) &= y_{N-1}(x+h)y_N(x+1/2h)T_N(x). \end{aligned}$$

For $i = 1, \dots, N-1$, the N -tuple $\mathbf{y}^{(i)} = (y_1, \dots, \tilde{y}_i, \dots, y_N)$ is called an immediate descendant of $\mathbf{y}$ in the direction i .

It follows from the N -th equation that the polynomial y_N belongs to a C_1 population with weight $y_{N-1}(x+h)T_N(x)$. Let $\tilde{y}_N$ be any element of that C_1 population. Then $\mathbf{y}^{(N)} = (y_1, \dots, y_{N-1}, \tilde{y}_N)$ is called an immediate descendant of $\mathbf{y}$ in the direction N .

For $i = 1, \dots, N$, immediate descendants $\mathbf{y}^{(i)}$ are fertile in the C_N sense.

From Lemma 3.3 we obtain that an N -tuple $\mathbf{y}$ represents a critical point of C_N type if and only if it is fertile and $\mathbf{y}^A$ is generic in sl_{2N+1} sense. If $\mathbf{y}$ represents a critical point of C_N type and if $(\mathbf{y}^{(i)})^A$ is generic in sl_{2N+1} sense, then $\mathbf{y}^{(i)}$ also represents an h -critical point of type C_N .

The C_N population originated at a critical point $\mathbf{y}$ is the minimal set $P(\mathbf{y})$ of fertile N -tuples such that $\mathbf{y} \in P(\mathbf{y})$ and if $\bar{\mathbf{y}} \in P(\mathbf{y})$ then the immediate descendants $\bar{\mathbf{y}}^{(i)} \in P(\mathbf{y})$.

Obviously, the C_N population is contained in the corresponding sl_{2N+1} population: if $\bar{\mathbf{y}} \in P(\mathbf{y})$ then $\bar{\mathbf{y}}^A \in P_A(\mathbf{y}^A)$, where we denote $P_A(\mathbf{y}^A)$ the sl_{2N+1} population of critical points originated at $\mathbf{y}^A$. Moreover, the fundamental space of the sl_{2N+1} population $P_A(\mathbf{y}^A)$ contains a flag with the property (5.3) and therefore is h -selfdual. We call this space the *fundamental space of the C_N population $P(\mathbf{y})$* .

The corresponding fundamental operator is the difference operator of order $2N+1$ obtained from (4.2) by substituting (7.1) and (5.2):

$$\begin{aligned} & \prod_{i=1}^{N-1} \left(\partial - \frac{y_{N-i}(x+(i+3/2)h)}{y_{N-i}(x+(i+1/2)h)} \frac{y_{N+1-i}(x+(i-1/2)h)}{y_{N+1-i}(x+(i+1/2)h)} T(x+ih) \times \right. \\ & \left. \prod_{s=1}^{i-1} \frac{T_{N-s}(x+(i-1/2)h)}{T_{N-s}(x+(i-3/2)h)} \right) \left(\partial - \frac{y_N(x+3/2h)}{y_N(x+1/2h)} \frac{y_N(x)}{y_N(x+h)} T(x) \right) \times \\ & \prod_{i=1}^{1 \rightarrow N} \left(\partial - \frac{y_{N+1-i}(x+h)}{y_{N+1-i}(x)} \frac{y_{N-i}(x)}{y_{N-i}(x+h)} \prod_{s=1}^{N-i} \frac{T_s(x+(N-i-s+1)h)}{T_s(x+(N-i-s)h)} \right), \end{aligned}$$

where $T(x) = \prod_{s=1}^N T_s(x+(N-s+1)h)/T_s(x+(N-s)h)$.

Clearly two populations of h -critical points of type C_N either do not intersect or coincide.

7.4. A C_N population and the B_N flag variety. Let $\mathbf{y}$ be an h -critical point of C_N type and let V be the fundamental space of the population $P_A(\mathbf{y}^A)$. Then V is an h -selfdual space of dimension $2N+1$ and therefore has a non-degenerate symmetric canonical bilinear form.

The special orthogonal Lie algebra of V consists of all traceless endomorphisms x of V such that $(xv, v') + (v, xv') = 0$ for all $v, v' \in V$. Let $\mathbf{u} = (u_1, \dots, u_{2N+1})$ be a Witt basis in V . We have $(u_i, u_{2N+2-i}) = (-1)^{i+1}$, $i = 1, \dots, N+1$, and $(u_i, u_j) = 0$ if $i + j \neq 2N+2$. The choice of the basis identifies the special orthogonal Lie algebra with a Lie subalgebra of sl_{2N+1} , which is denoted so_{2N+1} . The Lie algebra so_{2N+1} has the root system of type B_k .

The lower triangular part of so_{2N+1} is spanned by matrices $E_{i,i+1} + E_{2N+1-i, 2N+2-i}$ for $i = 1, \dots, N$. Denote these matrices $X_1, \dots, X_N$, respectively.

Now we describe the action of one-parametric subgroups in the special orthogonal group in the directions of X_i on the basis $\mathbf{u} = (u_1, \dots, u_{2N+1})$. We have

$$\begin{aligned} e^{cX_i} \mathbf{u} &= \\ &= (u_1, \dots, u_{i-1}, u_i + cu_{i+1}, u_{i+1}, \dots, u_{2N+1-i}, u_{2N+2-i} + cu_{2N+3-i}, \dots, u_{2N+1}), \\ e^{cX_i} \mathbf{u} &= (u_1, \dots, u_{N-1}, u_N + cu_{N+1} + c^2 u_{N+2}/2, u_{N+1} + cu_{N+2}, u_{N+2}, \dots, u_{2N+1}), \end{aligned}$$

where $i = 1, \dots, N-1$.

We also set

$$\begin{aligned} e^{\infty X_i} \mathbf{u} &= (u_1, \dots, u_{i-1}, u_{i+1}, u_i, \dots, u_{2N+1-i}, u_{2N+3-i}, u_{2N+2-i}, \dots, u_{2N+1}), \\ e^{\infty X_i} \mathbf{u} &= (u_1, \dots, u_{N-1}, u_{N+2}, u_{N+1}, u_N, \dots, u_{2N+1}). \end{aligned}$$

Note $e^{cX_i} \mathbf{u}$ is a Witt basis for all i, c and the corresponding flag is isotropic.

Denote $FL^\perp(V) \subset FL(V)$ the subvariety of all isotropic flags and define the C_N generating morphism $\beta : FL^\perp(V) \rightarrow \mathbf{P}(\mathbb{C}[x])^N, \mathcal{F}(\mathbf{u}) \mapsto (y_1^\mathcal{F}, \dots, y_N^\mathcal{F})$, where $y_i^\mathcal{F} = W^\dagger(u_1, \dots, u_i)$.

Lemma 7.2. *The set $\beta(\mathcal{F}(e^{cX_i} \mathbf{u}))$, $c \in \bar{\mathbb{C}}$ coincides with the set of all immediate descendants of $\beta(\mathcal{F}(\mathbf{u}))$ in the direction i .*

Proof. Follows from the Wronskian identities

$$\begin{aligned} W(y_i^\mathcal{F}, W^\dagger(u_1, \dots, u_{i-1}, u_{i+1}))(x) &= T_i(x) y_{i-1}^\mathcal{F}(x+h) y_{i+1}^\mathcal{F}(x), \\ W(y_N^\mathcal{F}, W^\dagger(u_1, \dots, u_{N-1}, u_{N+1}))(x) &= T_N(x) y_{N-1}^\mathcal{F}(x+h) y_N^\mathcal{F}(x+h/2), \\ W(y_N^\mathcal{F}, W^\dagger(u_1, \dots, u_{N-1}, u_{N+2}))(x) &= \\ &= T_N(x) y_{N-1}^\mathcal{F}(x+h) W^\dagger(u_1, \dots, u_{N-1}, u_{N+1})(x+h/2), \end{aligned}$$

where $i = 1, \dots, N-1$. □

Theorem 7.3. *The generating morphism β of type C_N gives rise to the isomorphism $FL^\perp(V) \rightarrow P(\mathbf{y}) \subset \mathbf{P}(\mathbb{C}[x])^N$.*

Proof. The generating morphism β is an isomorphism of $FL^\perp(V)$ to the image by Theorem 4.16. Clearly, the image of β contains the population $P(\mathbf{y})$. The image of β coincides with $P(\mathbf{y})$ by Lemma 7.2. □

Note that $FL^\perp$ is isomorphic to the flag variety of the special symplectic group which corresponds to the B_N root system. This flag variety has a Bruhat cell decomposition $FL^\perp(V) = \sqcup_{w \in \mathcal{W}} G_w^{\mathcal{F}(\infty), B}$, where $\mathcal{W}$ is the B_N Weyl group. The Weyl groups of B_N and C_N are canonically identified. See for details, e.g. [MV1], section 7.4.

As before, the set of all N -tuples in a C_N population of the same degree is the Bruhat cell in the B_N flag variety.

Theorem 7.4. *The set of N -tuples $\bar{\mathbf{y}}$ in a C_N population $P(\mathbf{y})$ associated with a weight at infinity $w \cdot \Lambda_\infty$, where Λ_∞ is integral dominant, coincides with the image of Bruhat cell $\beta(G_w^{\mathcal{F}(\infty), B})$.*

Proof. Completely parallel to the proof of Corollary 7.14 in [MV1]. $\square$

8. APPENDIX A: AN EXAMPLE OF AN sl_3 POPULATION

Consider the population of h -critical points associated to $N = 2$ and $n = 0$ and originated at $\mathbf{y}^0 = (1, 1)$. The pair $(1, 1)$ represents the h -critical point with no variables. We claim that this population consists of pairs of non-zero polynomials $\mathbf{y} = (y_1, y_2)$, where

$$y_i = a_{2,i}x^2 + a_{1,i}x + a_{0,i}, \quad i = 1, 2, \quad (8.1)$$

and

$$(a_{1,1} + a_{2,1}h)(a_{1,2} - a_{2,2}h) = 2a_{0,1}a_{2,2} + 2a_{2,1}a_{0,2}.$$

For any generic pair $\mathbf{y} = (y_1, y_2)$ of this form, the roots of the polynomials y_1, y_2 form an h -critical point with the initial data where $n = 0$. In other words, the roots of y_1, y_2 satisfy the equations

$$\prod_{k \neq j} \frac{t_j^{(1)} - t_k^{(1)} - h}{t_j^{(1)} - t_k^{(1)} + h} \prod_{k=1}^{l_2} \frac{t_j^{(1)} - t_k^{(2)}}{t_j^{(1)} - t_k^{(2)} - h} = 1, \quad j = 1, \dots, l_1,$$

$$\prod_{k=1}^{l_1} \frac{t_j^{(2)} - t_k^{(1)} + h}{t_j^{(2)} - t_k^{(1)}} \prod_{k \neq j} \frac{t_j^{(2)} - t_k^{(2)} - h}{t_j^{(2)} - t_k^{(2)} + h} = 1, \quad j = 1, \dots, l_2,$$

where $l_1 = \deg y_1$ and $l_2 = \deg y_2$.

Equations (3.6) take the form

$$W(y_1, \tilde{y}_1)(x) = y_2(x), \quad W(y_2, \tilde{y}_2)(x) = y_1(x + h), \quad (8.2)$$

and the reproduction procedure works as follows. We start with $\mathbf{y}^0 = (1, 1)$. Equations (8.2) have the form $W(1, \tilde{y}_1)(x) = 1$, $W(1, \tilde{y}_2)(x) = 1$. Using the second of them, we get pairs $\mathbf{y} = (1, x + a)$ for all numbers a . Equations (8.2) now are $W(1, \tilde{y}_1) = x + a$, $W(x + a, \tilde{y}_2) = 1$. Using the first equation we get pairs $\mathbf{y} = (x^2 + (2a - h)x + b, x + a)$ for all a, b . Equations (8.2) take the form $W(x^2 + (2a - h)x + b, \tilde{y}_1) = x + a$, $W(x + a, \tilde{y}_2) = (x + h)^2 + (2a - h)(x + h) + b$. Using the second of them we get $\mathbf{y} = (x^2 + (2a - h)x + b, x^2 + cx + ac - ah - b)$ for all a, b, c . It is easy to see that the

union of all those pairs is our population, and nothing else can be constructed starting from $\mathbf{y}^0 = (1, 1)$.

Now, it is easy to see that the family of pairs (8.1) (where each pair is considered up to multiplication of its coordinates by non-zero numbers) is isomorphic as an algebraic variety to the variety of all full flags in the three dimensional vector space $V = \mathbb{C}_2[x]$ of the first coordinates of the pairs. Namely, y_1 generates a line in V and y_2 defines a plane in V containing the line generated by y_1 . The space V is the fundamental space of the population.

In the example above the possible degrees of polynomials y_1, y_2 are $(0,0), (1,0), (0,1), (1,2), (2,1), (2,2)$. The corresponding parts of the family are isomorphic to open Bruhat cells of dimensions 0, 1, 1, 2, 2, 3, respectively.

The fundamental space of the population V is the space of polynomials of degree at most 2. This is an h -selfdual space. The canonical form in the basis $(1, x, x(x-1)/2)$ is described by the matrix:

$$\begin{pmatrix} 0 & 0 & 1 \\ 0 & -1 & 1/2 \\ 1 & 1/2 & -1/8 \end{pmatrix}.$$

It is a symmetric non-degenerate form.

The basis $(1, x - 1/2, (x^2 - x - 1/8)/2)$ is a Witt basis.

The set of isotropic vectors is the C_1 population. In this example, this is the set of polynomials of the form

$$(x(x-1) - 1/8)/2 + \alpha(x - 1/2) + \alpha^2/2 = (x + \alpha - 1/2)^2 - 1/8,$$

where $\alpha \in \mathbb{C}$ and also $1 \in \mathcal{V}$ which corresponds to $\alpha = \infty$.

9. APPENDIX B: THE WRONSKIAN IDENTITIES

In this appendix we collect identities involving discrete Wronskians. All functions in this section are functions of one variable x .

Set

$$\Delta f(x) = f(x+h) - f(x).$$

Set $\Delta^{(0)}f(x) = f(x)$ and

$$\Delta^{(n+1)}f(x) = \Delta(\Delta^{(n)}f)(x).$$

The discrete Wronskian of functions $g_1, \dots, g_s$ is defined by

$$W(g_1, \dots, g_s)(x) = \det(g_i(x + (j-1)h))_{i,j=1}^s = \det(\Delta^{(j-1)}g_i(x))_{i,j=1}^s.$$

We follow the convention that for $s = 0$ the corresponding discrete Wronskian equals 1. In this section we write $W_s(g_1, \dots, g_s)$ instead of $W(g_1, \dots, g_s)$ to stress the order of the Wronskian.

Lemma 9.1. *We have $W_{s+1}(1, g_1, \dots, g_s)(x) = W_s(\Delta g_1, \dots, \Delta g_s)(x)$.* □

Now we describe the Wronskian $W_s(\Delta g_1, \dots, \Delta g_s)(x)$ in more detail. Apriori, this Wronskian consists of $2^s s!$ terms. However, there are cancellations. We describe the surviving terms.

Lemma 9.2. *The Wronskian $W_s(\Delta g_1, \dots, \Delta g_s)(x)$ is equal to the alternating sum of $(s+1)!$ terms of the form $\prod_{i=1}^s g_i(x + \omega_i h)$ where $(\omega_1, \dots, \omega_s)$ is a permutation ω of the set $\{0, \dots, \hat{j}, \dots, s\}$ for some j . The sign of this term is $(-1)^j \text{sgn}(\omega)$.*

Proof. This lemma is straightforward. $\square$

Now we proceed to other identities.

Lemma 9.3. *We have $W_s(fg_1, \dots, fg_s)(x) = W_s(g_1, \dots, g_s)(x) \prod_{j=0}^{s-1} f(x + jh)$.* $\square$

For given functions $g_1, \dots, g_{s+1}$ and an integer k , $0 \leq k \leq s$, denote $V_{s-k+1}(i) = W_{s-k+1}(g_1, \dots, g_{s-k}, g_i)$.

Lemma 9.4. *We have*

$$W_{k+1}(V_{s-k+1}(s-k+1), \dots, V_{s-k+1}(s+1))(x) = \\ W_{s+1}(g_1, \dots, g_{s+1})(x) \prod_{j=1}^k W_{s-k}(g_1, \dots, g_{s-k})(x + jh) .$$

Proof. This lemma is proved by induction on s . The case of $k = s$ is trivial. Suppose that the lemma is proved for $s = k, k+1, \dots, s_0 - 1$. Divide both sides of the identity for $s = s_0$ by $\prod_{j=0}^{s_0-k} \prod_{i=0}^k g_1(x + (i+j)h)$ and use Lemma 9.3 to carry g_1 inside all Wronskians. Then one of the functions in each Wronskian is 1 and we can reduce the order by Lemma 9.1. Then the identity for $s = s_0$ follows from the induction hypothesis applied to $f_i = \Delta(g_{i+1}/g_1)$, $i = 1, \dots, s_0$. $\square$

For given functions $g_1, \dots, g_{s+1}$, denote $W_s(i) = W(g_1, \dots, \hat{g}_i, \dots, g_{s+1})$ the Wronskian of all functions except g_i .

Lemma 9.5. *We have*

$$W_{k+1}(W_s(s+1), W_s(s), \dots, W_s(s-k+1))(x) = \\ W_{s-k}(g_1, \dots, g_{s-k})(x + kh) \prod_{j=1}^k W_{s+1}(g_1, \dots, g_{s+1})(x + (j-1)h) .$$

Proof. First we prove the case of $s = k$ by induction on k . The case of $k = 0$ is trivial. Suppose the case of $k < k_0$ is proved. Divide both sides of our identity in the case of $s = k = k_0$ by $\prod_{i=0}^{k_0-1} \prod_{j=0}^{k_0} g_1(x + (i+j)h)$. By Lemmas 9.1 and 9.3 we are reduced to the identity

$$W_{k_0+1}(W_{k_0-1}^{\Delta h}[k_0], \dots, W_{k_0-1}^{\Delta h}[1], W_{k_0}(h_1, \dots, h_{k_0})) = \prod_{i=0}^{k_0-1} W(\Delta h_1, \dots, \Delta h_{k_0})(x + ih),$$

where $h_i = g_{i+1}/g_1$ and $W_{k_0-1}^{\Delta h}[i] = W_{k_0-1}(\Delta h_1, \dots, \widehat{\Delta h_i}, \dots, \Delta h_{k_0})$.

The left hand side of the last identity is a determinant of size $k_0 + 1$. Add to the last row the row number i with coefficient $(-1)^{k_0-i+1} h_{k_0-i+1}(x + k_0 h)$, $i = 1, \dots, k_0$. Then using Lemma 9.2 we observe that the last row becomes

$$(0, \dots, 0, W_{k_0}(\Delta h_1, \dots, \Delta h_{k_0})(x + (k_0 - 1)h))$$

and the lemma for $k = k_0$ follows from the induction hypothesis applied to functions $f_1 = \Delta h_1, \dots, f_{k_0} = \Delta h_{k_0}$.

Now we continue by induction on s . Suppose that the lemma is proved for $s = k, \dots, s_0 - 1$. Divide both sides of the identity for $s = s_0$ by $\prod_{i=0}^{s_0-1} \prod_{j=0}^k g_1(x + (i+j)h)$. Then the identity for $s = s_0$ follows from the induction hypothesis applied to $f_i = \Delta(g_{i+1}/g_1)$, $i = 0, \dots, s_0 - 1$. $\square$

REFERENCES

- [BIK] N.M. Bogoliubov, A.G. Izergin and V.E. Korepin, *Quantum Inverse Scattering Method and Correlation Functions*, Cambridge University Press, Cambridge, 1993.
- [F] W. Fulton, *Intersection Theory*, Springer-Verlag, 1984.
- [Fd] L.D. Faddeev, *Lectures on quantum inverse scattering method*, in *Integrable Systems*, Nankai Lectures on Math Phys., 1987 (X-C. Song. ed.), World Scientific, Singapore (1990), 23-70.
- [FT] L.D. Faddeev, L.A. Takhtajan, *Quantum inverse problem method and the Heisenberg XYZ model*, Russian Math. Survey, 34 (1979), **5**, 11-68.
- [GH] Ph. Griffiths, J. Harris, *Principles of algebraic geometry*, A Wiley-Interscience Publication, 1994.
- [MV1] E. Mukhin and A. Varchenko, *Critical Points of Master Functions and Flag Varieties*, math.QA/0209017 (2002), 1-49.
- [MV2] E. Mukhin and A. Varchenko, *The quantized Knizhnik-Zamolodchikov equation in tensor products of irreducible $sl(2)$ -modules*, Calogero-Moser-Sutherland Models workshop, Springer (2000), 347-384.
- [TV1] V. Tarasov and A. Varchenko, *Geometry of q -hypergeometric functions as a bridge between Yangians and quantum affine algebras*, Invent. math. 128, (1997) 501-588.
- [TV2] V. Tarasov and A. Varchenko, *Completeness of Bethe vectors and Difference equations with Regular Singular points*, International Mathematics Research Notices (1995), **no. 13**, 637-669.

E.M.: DEPARTMENT OF MATHEMATICAL SCIENCES, INDIANA UNIVERSITY - PURDUE UNIVERSITY INDIANAPOLIS, 402 NORTH BLACKFORD ST, INDIANAPOLIS, IN 46202-3216, USA,
MUKHIN@MATH.IUPUI.EDU

A.V.: DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NORTH CAROLINA AT CHAPEL HILL, CHAPEL HILL, NC 27599-3250, USA, ANV@EMAIL.UNC.EDU